

# Applied/Numerical Analysis Qualifying Exam

January 11, 2011

## Cover Sheet – Part I

**Policy on misprints:** The qualifying exam committee tries to proofread exams as carefully as possible. Nevertheless, the exam may contain a few misprints. If you are convinced a problem has been stated incorrectly, indicate your interpretation in writing your answer. In such cases, do *not* interpret the problem so that it becomes trivial.

Name \_\_\_\_\_

## Part 1: Applied Analysis

**Instructions:** Do any 3 of the 4 problems in this part of the exam. Show all of your work clearly. Please indicate which of the 4 problems you are skipping.

- (1) Given  $w \in C[0, 1]$ , with  $w(x) > 0$  on  $[0, 1]$ , let  $L_w^2[0, 1]$  be the weighted Hilbert space with the inner product

$$\langle f, g \rangle_w = \int_0^1 f(x) \overline{g(x)} w(x) dx,$$

where  $f, g$  are in  $L^2[0, 1]$ . In addition, let  $\{\phi_n(x)\}_{n=0}^\infty$  be the set of orthogonal polynomials generated by using the Gram-Schmidt process on  $\{1, x, x^2, \dots\}$  in the inner product for  $L_w^2$ . Assume that  $\phi_n(x) = x^n + \text{lower powers}$ .

- (a) State the Weierstrass Approximation Theorem and briefly sketch its proof. (Use no more than a page or so.)
- (b) You are given that  $C[0, 1]$  is dense in  $L^2[0, 1]$ . Show that the orthogonal polynomials  $\{\phi_n(x)\}_{n=0}^\infty$  form a complete, orthogonal set in  $L_w^2[0, 1]$ .
- (2) Consider the differential operator  $Lu(x) = -((x+1)u)'$ , with  $x \in [0, 1]$ .
- (a) Show that if  $D(L) := \{u \in L^2 \mid Lu \in L^2 \text{ and } u(0) = 0 = u'(1)\}$ , then  $L$  is self adjoint and positive definite.
- (b) Find the Green's function for  $L$  having the domain  $D(L)$  above.
- (c) Briefly explain why the eigenfunctions this operator are complete in  $L^2[0, 1]$ .
- (3) In the problem below, use the Fourier transform conventions

$$\begin{aligned} \mathcal{F}[f](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx \\ \mathcal{F}^{-1}[\hat{f}](x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(\omega) e^{i\omega x} d\omega. \end{aligned}$$

As usual,  $\hat{f} = \mathcal{F}[f]$ .

- (a) Show  $\mathcal{F}^4 = I$ . (Hint:  $\mathcal{F}[f(x)] = \mathcal{F}^{-1}[f(-x)]$ .)
- (b) You are given that the equation  $-u_n'' + x^2 u_n = (2n+1)u_n$  has, up to a constant multiple, a unique solution  $u_n \in L^2(\mathbb{R})$ , for  $n = 0, 1, \dots$ . (You may assume that the solution is smooth enough and decays fast enough to be in Schwartz space.) Show that  $u_n$  is an eigenfunction of the Fourier transform; that is,  $\hat{u}_n(\omega) = \lambda_n u_n(\omega)$ . Also, show that  $\lambda_n^4 = 1$ .
- (4) Let  $k(x, y) = x^4 y^{12}$  and consider the operator  $Ku(x) = \int_0^1 k(x, y) u(y) dy$ .
- (a) Show that  $K$  is a Hilbert-Schmidt operator and that  $\|K\|_{\text{op}} \leq \frac{1}{10}$ .
- (b) State the Fredholm Alternative for the operator  $L = I - \lambda K$ . Explain why it applies in this case. Find all values of  $\lambda$  such that  $Lu = f$  has a unique solution for all  $f \in L^2[0, 1]$ .
- (c) Use a Neumann series to find the resolvent  $(I - \lambda K)^{-1}$  for  $\lambda$  small. Sum the series to find the resolvent.

# Applied/Numerical Analysis Qualifying Exam

January 11, 2011

## Cover Sheet – Part II

**Policy on misprints:** The qualifying exam committee tries to proofread exams as carefully as possible. Nevertheless, the exam may contain a few misprints. If you are convinced a problem has been stated incorrectly, indicate your interpretation in writing your answer. In such cases, do *not* interpret the problem so that it becomes trivial.

Name \_\_\_\_\_

## Part 2: Numerical Analysis

**Instructions:** Do all problems in this part of the exam. Show all of your work clearly.

**Problem 1:** Consider the following two-points boundary value second order problem in 1-D: Find a function  $u$  defined a.e. in  $]0, 1[$  such that

$$(1) \quad \begin{aligned} & -(xK(x)u'(x))' + xq(x)u(x) = xf(x) \text{ a.e. in } ]0, 1[, \\ & \lim_{x \rightarrow 0} (xu'(x)) = 0 \text{ and } K(1)u'(1) + u(1) = 0, \end{aligned}$$

where  $K \in \mathcal{C}^1([0, 1])$ ,  $q \in \mathcal{C}^0([0, 1])$  and  $f \in L^2(0, 1)$  are given functions. Assume that there exists a constant  $\kappa_0 > 0$  such that  $K(x) \geq \kappa_0$  and  $q(x) \geq 0$  for all  $x \in [0, 1]$ . Let

$$V = \{v \in L^2_{\text{loc}}(0, 1); \sqrt{x}v \in L^2(0, 1), \sqrt{x}v' \in L^2(0, 1)\}.$$

Accept as a fact that  $V$  is a Hilbert space for the norm

$$\|v\|_V = \left( \|\sqrt{x}v\|_{L^2(0,1)}^2 + \|\sqrt{x}v'\|_{L^2(0,1)}^2 \right)^{1/2},$$

and  $\mathcal{C}^1([0, 1])$  is dense in  $V$  for this norm.

- (1) Derive the variational formulation (also called weak formulation) of problem (1) in the space  $V$ .
- (2) Prove that the corresponding bilinear form of this variational formulation is elliptic (or coercive) in  $V$ .

**Hint.** First show that all functions  $v$  of  $\mathcal{C}^1([0, 1])$  satisfy

$$\int_0^1 v(x)^2 dx = v^2(1) - 2 \int_0^1 xv(x)v'(x) dx$$

and then establish the following variant of Poincaré's inequality

$$\forall v \in V, \|\sqrt{x}v\|_{L^2(0,1)} \leq \alpha \left( v^2(1) + \|\sqrt{x}v'\|_{L^2(0,1)}^2 \right)^{\frac{1}{2}}$$

for some constant  $\alpha > 0$ . Based on this equality deduce the ellipticity.

- (3) Choose an integer  $N \geq 2$ , set  $h = 1/N$ , let  $x_i = ih$ ,  $0 \leq i \leq N$  and define the finite element space

$$V_h = \{v_h \in \mathcal{C}^0([0, 1]); v_h|_{]x_i, x_{i+1}[} \in \mathcal{P}_1, 0 \leq i \leq N-1\}.$$

Show that  $V_h$  is a subspace of  $V$ . Discretize the variational problem in this space. Prove existence and uniqueness of the discrete solution and establish an error estimate without estimating the norms of the interpolation errors.

**Problem 2:** Let  $\Omega$  be a bounded domain in  $\mathbb{R}^2$  with polygonal boundary  $\partial\Omega$ . Let

$$H_0^1(\Omega) = \{v \in H^1(\Omega) : v(x) = 0 \forall x \in \partial\Omega\}$$

be the standard Sobolev space of functions defined on  $\Omega$  that vanish on the boundary.

In all that follows,  $T > 0$  is a given final time,  $c > 0$  is a constant, and  $u_0 \in \mathcal{C}^0(\Omega)$  are given functions. Consider the parabolic equation: Find a function  $u$  defined a.e. in  $\Omega \times ]0, T[$  solution of

$$(2) \quad \begin{aligned} & \frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x_1^2} - \frac{\partial^2 u}{\partial x_2^2} + cu = 0 \text{ a.e. in } \Omega \times ]0, T[, \\ & u(x, t) = 0 \text{ a.e. in } \partial\Omega \times ]0, T[, \\ & u(x, 0) = u_0(x) \text{ a.e. in } \Omega. \end{aligned}$$

Accept as a fact that problem (2) has one and only one solution  $u$  in  $L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ .

Let  $\mathcal{T}_h$  be a finite element partition of  $\Omega$  into triangles  $\tau$  of diameter  $h_\tau \leq h$ . Further, let

$$W_h = \{v_h \in C^0(\bar{\Omega}); \forall \tau \in \mathcal{T}_h, v_h|_\tau \in \mathcal{P}_1, v_h|_{\partial\Omega} = 0\},$$

be a finite element space of continuous piece-wise linear functions over  $\mathcal{T}_h$ .

Consider the fully discrete backward Euler implicit approximation of (2): for  $K$  a positive integer, set  $k = T/K$ , define  $t_n = nk$ ,  $0 \leq n \leq K$ , and for each  $0 \leq n \leq K - 1$ , knowing  $u_h^n \in W_h$  find  $u_h^{n+1} \in W_h$  such that

$$(3) \quad \forall v_h \in W_h, \frac{1}{k}(u_h^{n+1} - u_h^n, v_h) + a(u_h^{n+1}, v_h) = 0, \quad n = 0, 1, \dots, K, \quad u_h^0 = I_h(u_0).$$

Here  $(\cdot, \cdot)$  is the inner product in  $L^2(\Omega)$ , the bilinear form  $a(u_h^{n+1}, v_h)$  comes from the variational formulation of problem (2), and  $I_h$  is the Lagrange interpolation operator in  $W_h$ . Write the expression of  $a(u_h^{n+1}, v_h)$ .

(1) Show that (3) defines a unique function  $u_h^{n+1}$  in  $W_h$ .

(2) Prove the following stability estimate

$$(4) \quad \sup_{1 \leq n \leq K} \|u_h^n\|_{L^2(\Omega)}^2 + k \sum_{n=1}^K |u_h^n|_{H^1(\Omega)}^2 \leq \|u_h^0\|_{L^2(\Omega)}^2.$$

(3) Also prove the estimate

$$(5) \quad \sup_{1 \leq n \leq K} |u_h^n|_{H^1(\Omega)} \leq |u_h^0|_{H^1(\Omega)}.$$

**Problem 3:** Consider the interval  $(0, 1)$  and the set of continuous functions  $\hat{v}$  defined on  $[0, 1]$ . Let  $\hat{a}_1 = 0$ ,  $\hat{a}_2 = \frac{1}{2}$ ,  $\hat{a}_3 = 1$ .

(1) Consider the following two sets of degrees of freedom

$$\Sigma_1 = \{\hat{v}(\hat{a}_j), j = 1, 2, 3\} \quad \Sigma_2 = \{\hat{v}(\hat{a}_1), \hat{v}(\hat{a}_3), \int_0^1 \hat{v}(s) ds\}.$$

Write down the basis functions of  $\mathcal{P}_2$  (for both sets of degrees of freedom) such that

(a)  $p_i \in \mathcal{P}_2$ ,  $1 \leq i \leq 3$ , satisfying:  $p_i(\hat{a}_j) = \delta_{i,j}$ ,  $1 \leq i, j \leq 3$  for the set  $\Sigma_1$ ;

(b)  $q_i \in \mathcal{P}_2$ ,  $1 \leq i \leq 3$ , satisfying:

$$q_i(\hat{a}_j) = \delta_{i,j}, \int_0^1 q_i(s) ds = 0, \quad i = 1, 3, j = 1, 3,$$

$$\int_0^1 q_2(s) ds = 1, q_2(\hat{a}_1) = q_2(\hat{a}_3) = 0, \quad \text{for the set } \Sigma_2.$$

In both cases, write down the FE interpolant  $\hat{\Pi}(\hat{w})$  of a given function  $\hat{w} \in C^0([0, 1])$ .

(2) Consider the interval  $[a, b]$ , let  $F$  map  $[0, 1]$  onto  $[a, b]$ , and let  $v$  be given in  $H^3(a, b)$ .

Define  $\Pi(v)$  by  $(\Pi(v)) \circ F = \hat{\Pi}(v \circ F)$ . Give the Bramble Hilbert argument to get an estimate in terms of  $h = b - a$  for the error

$$\|v' - \Pi(v)'\|_{L^2(a,b)}.$$

Explain how to modify the proof when  $v$  is less regular, e.g  $v \in H^2(a, b)$ .