

Quiz 2 (Notes, books, and calculators are not authorized)

Show all your work in the blank space you are given on the exam sheet. Always justify your answer. Answers with **no justification will not be graded**.

**Question 1:** Let  $\phi$  be a non-zero solution to the eigenvalue problem  $-\partial_{xx}\phi(x) = \lambda\phi(x)$ ,  $x \in (0, \pi)$ ,  $\phi(0) = 0$ ,  $\phi(\pi) = 0$ . Determine the sign of  $\lambda$  using the energy method.

Multiply the equation by  $\phi$ , integrate over  $(0, \pi)$ , and apply the fundamental theorem of calculus (i.e. integrate by parts):

$$\begin{aligned} -\int_0^\pi \phi(x)\partial_{xx}\phi(x)dx &= -\int_0^\pi (\partial_x(\phi(x)\partial_x\phi(x)) - (\partial_x\phi(x))^2)dx \\ &= -\phi(\pi)\partial_x\phi(\pi) + \phi(0)\partial_x\phi(0) + \int_0^\pi (\partial_x\phi(x))^2dx = \lambda \int_0^\pi (\phi(x))^2dx. \end{aligned}$$

In conclusion

$$\int_0^\pi (\partial_x\phi(x))^2dx = \lambda \int_0^\pi (\phi(x))^2dx.$$

Assuming that  $\phi$  is nonzero, we obtain that  $\lambda = \int_0^\pi (\partial_x\phi(x))^2dx / \int_0^\pi (\phi(x))^2dx \geq 0$ , i.e.  $\lambda$  is non-negative. If  $\lambda = 0$  then  $\partial_x\phi = 0$ , which implies that  $\phi$  is constant. The boundary conditions imply that  $\phi = 0$  which contradicts our assumption that  $\phi$  is non-zero. In conclusion  $\lambda$  is negative.

**Question 2:** Compute all the positive eigenvalues  $\lambda$  to the above eigenvalue problem.

Owing to  $\lambda$  being non-negative, the general solution to  $-\partial_{xx}\phi(x) = \lambda\phi(x)$  is  $\phi(x) = a \cos(\sqrt{\lambda}x) + b \sin(\sqrt{\lambda}x)$ . The boundary conditions imply that  $\phi(0) = 0 = a$  and  $\phi(\pi) = b \sin(\sqrt{\lambda}\pi) = 0$ . As a result,  $\sqrt{\lambda}\pi = n\pi$ , where  $n \in \mathbb{N}$ , i.e.

$$\lambda = n^2, \quad n \in \mathbb{N}.$$

**Question 3:** Consider the heat equation  $\partial_t u(x, t) - 3\partial_{xx}u(x, t) = 0$ ,  $\partial_x u(0, t) = 0$ ,  $\partial_x u(1, t) = 0$ ,  $u(x, 0) = u_0(x)$ ,  $t > 0$ ,  $x \in (0, 1)$ . The general solution is  $u(x, t) = \sum_{n=0}^{\infty} A_n \cos(n\pi x)e^{-3n^2\pi^2 t}$ . Compute the solution corresponding to the initial data  $u_0(x) = 5 \cos(4\pi x)$ .

The solution contains one term only, corresponding to  $n = 4$ ,

$$u(x, t) = 5 \cos(4\pi x)e^{-48\pi^2 t}.$$

**Question 4:** Assume that the following equation has a smooth solution:  $-\partial_x((1+x^2)\partial_x T(x)) + \partial_x T(x) + T(x) = 2x - 1$ ,  $T(a) = 1$ ,  $T(b) = \pi$ ,  $x \in [a, b]$ ,  $t > 0$ , where  $k > 0$ . Prove that this solution is unique by using the energy method. (Hint: Do not try to simplify  $-\partial_x((1+x^2)\partial_x T)$ .)

Assume that there are two solutions  $T_1$  and  $T_2$ . Let  $\phi = T_2 - T_1$ . Then

$$-\partial_x((1+x^2)\partial_x \phi(x)) + \partial_x \phi(x) + \phi(x) = 0, \quad \phi(a) = 0, \quad \phi(b) = 0$$

Multiply the PDE by  $\phi$ , integrate over  $(a, b)$ , and integrate by parts (i.e. apply the fundamental theorem of calculus):

$$\begin{aligned} 0 &= \int_a^b (-\partial_x((1+x^2)\partial_x \phi(x))\phi(x) + (\partial_x \phi(x))\phi(x) + (\phi(x))^2) dx \\ &= \int_a^b (-\partial_x(\phi(x)(1+x^2)\partial_x \phi(x)) + (1+x^2)(\partial_x \phi(x))^2 + \partial_x(\frac{1}{2}\phi(x)^2) + (\phi(x))^2) dx \\ &= \int_a^b ((1+x^2)(\partial_x \phi(x))^2 + (\phi(x))^2) dx \end{aligned}$$

This implies  $\int_a^b (\phi(x))^2 dx = 0$ , i.e.  $\phi = 0$ , meaning that  $T_2 = T_1$ .