

CRITICAL POINTS OF DISCRETE PERIODIC OPERATORS

M. FAUST AND F. SOTTILE

ABSTRACT. We study the spectra of operators on periodic graphs using methods from combinatorial algebraic geometry. Our main result is a bound on the number of complex critical points of the Bloch variety, together with an effective criterion for when this bound is attained. We show that this criterion holds for \mathbb{Z}^2 - and \mathbb{Z}^3 -periodic graphs with sufficiently many edges and use our results to establish the spectral edges conjecture for some \mathbb{Z}^2 -periodic graphs.

INTRODUCTION

The spectrum of a \mathbb{Z}^d -periodic self-adjoint discrete operator L consists of intervals in \mathbb{R} . Floquet theory reveals that the spectrum is the image of the coordinate projection to \mathbb{R} of the *Bloch variety* (also known as the dispersion relation), an algebraic hypersurface in $(S^1)^d \times \mathbb{R}$. This coordinate projection defines a function λ on the Bloch variety, which is our main object of study.

When the operator is discrete, the complexification of the Bloch variety is an algebraic variety in $(\mathbb{C}^\times)^d \times \mathbb{C}$. Thus techniques from algebraic geometry and related areas may be used to address some questions in spectral theory. In the 1990's Gieseke, Knörrer, and Trubowitz [16] used algebraic geometry to study the Schrödinger operator on the square lattice \mathbb{Z}^2 with a periodic potential and established a number of results, including Floquet isospectrality, the irreducibility of its Fermi varieties, and determined the density of states. Recently there has been a surge of interest in using algebraic methods in spectral theory. This includes investigating the irreducibility of Bloch and Fermi varieties [11, 12, 23, 25], Fermi isospectrality [26], density of states [20], and extrema and critical points of the projection λ on Bloch varieties [1, 9, 25]. We use techniques from combinatorial algebraic geometry and geometric combinatorics [31] to study critical points of the function λ on the Bloch variety of a discrete periodic operator. We now discuss motivation and sketch our results. Some background on spectral theory is sketched in Section 1, and Section 2.1 gives some background from algebraic geometry.

An old and widely believed conjecture in mathematical physics concerns the structure of the Bloch variety near the edges of the spectral bands. Namely, that for a sufficiently general operator L (as defined in Section 1.1), the extrema of the band functions λ_j on the Bloch variety are nondegenerate in that their Hessians are nondegenerate quadratic forms. This

2010 *Mathematics Subject Classification.* 81U30, 81Q10, 14M25.

Key words and phrases. Bloch variety, Schrödinger operator, Kushnirenko Theorem, Toric variety, Newton polytope.

Research supported in part by Simons Collaboration Grant for Mathematicians 636314 and NSF grants DMS-2246031, DMS-2201005, DMS-2052572, and DMS-2000345.

spectral edges nondegeneracy conjecture is stated in [22, Conj. 5.25], and it also appears in [5, 21, 27, 28]. Important notions, such as effective mass in solid state physics, the Liouville property, Green’s function asymptotics, Anderson localization, homogenization, and many other assumed properties in physics, depend upon this conjecture.

The spectral edges conjecture states that for generic parameters, each extreme value is attained by a single band, the extrema are isolated, and the extrema are nondegenerate. We discuss progress for discrete operators on periodic graphs. In 2000, Klopp and Ralston [18] proved that for Laplacians with generic potential each extreme value is attained by a single band. In 2015, Filonov and Kachkovskiy [13] gave a class of two-dimensional operators for which the extrema are isolated. They also show [13, Sect. 6] that the spectral edges conjecture may fail for a Laplacian with general potential, which does not have generic parameters in the sense of Section 1.1. Most recently, Liu [25] proved that the extrema are isolated for the Schrödinger operator acting on the square lattice.

We consider a property which implies the spectral edges nondegeneracy conjecture: A family of operators has the *critical points property* if for almost all operators in the family, all critical points of the function λ (not just the extrema) are nondegenerate. Algebraic geometry was used in [9] to prove the following dichotomy: For a given algebraic family of discrete periodic operators, either the critical points property holds for that family, or almost all operators in the family have Bloch varieties with degenerate critical points.

In [9], this dichotomy was used to establish the critical points property for the family of Laplace-Beltrami difference operators on the \mathbb{Z}^2 -periodic diatomic graph of Figure 1. Bloch varieties for these operators were shown to have at most 32 critical points. A single example was computed to have 32 nondegenerate critical points. Standard arguments from algebraic geometry (see Section 5) implied that, for this family, the critical points property, and therefore also the spectral edges nondegeneracy conjecture, holds.

We extend part of that argument to operators on many periodic graphs. Let L be a discrete operator on a \mathbb{Z}^d -periodic graph Γ (see Section 1). Its (complexified) Bloch variety is a hypersurface in the product $(\mathbb{C}^\times)^d \times \mathbb{C}$ of a complex torus and the complex line defined by a Laurent polynomial $D(z, \lambda)$. The last coordinate λ , corresponding to projection onto the spectral axis, is the function on the Bloch variety whose critical points we study. Accordingly, we will call critical points of the function λ on the Bloch variety “critical points of the Bloch variety.” One contribution of this paper is to shift focus from spectral band functions λ_j defined on a compact torus to a global function on the complex Bloch variety. Another is to use the perspective of nonlinear optimization to address a question concerning the spectrum of a discrete periodic operator.

We state our first result. Let Γ be a connected \mathbb{Z}^d -periodic graph (as in Section 1.1). Fix a fundamental domain W for the \mathbb{Z}^d -action on the vertices of Γ . The support $\mathcal{A}(\Gamma)$ of Γ records the local connectivity between translates of the fundamental domain. It is the set of $a \in \mathbb{Z}^d$ such that Γ has an edge with endpoints in both W and $a+W$.

Theorem A. *The function λ on the Bloch variety of a discrete operator on Γ has at most*

$$d! |W|^{d+1} \text{vol}(\mathcal{A}(\Gamma))$$

isolated critical points. Here, $\text{vol}(\mathcal{A}(\Gamma))$ is the Euclidean volume of the convex hull of $\mathcal{A}(\Gamma)$.

This bound uses an outer approximation for the Newton polytope of $D(z, \lambda)$ (see Lemma 4.1) and a study of the equations defining critical points of the function λ on the Bloch variety, called the *critical point equations* (8). Corollary 2.5 is a strengthening of Theorem A. When the bound is attained all critical points are isolated.

Example. We illustrate Theorem A on the example from [9, Sect. 4]. Figure 1 shows a periodic graph Γ with $d = 2$ whose fundamental domain W has two vertices and its support




FIGURE 1. A dense periodic graph Γ with the convex hull of $\mathcal{A}(\Gamma)$.

$\mathcal{A}(\Gamma)$ consists of the columns of the matrix $\begin{pmatrix} 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & -1 \end{pmatrix}$. Figure 1 also displays the convex hull of $\mathcal{A}(\Gamma)$. As $|W| = 2$ and $\text{vol}(\mathcal{A}(\Gamma)) = 2$, Theorem A implies that any Bloch variety for an operator on Γ has at most

$$d! |W|^{d+1} \text{vol}(\mathcal{A}(\Gamma)) = 2! \cdot 2^{2+1} \cdot 2 = 32$$

critical points, which is the bound demonstrated in [9]. ◇

The bound of Corollary 2.5 arises as follows. There is a natural compactification of $(\mathbb{C}^\times)^d \times \mathbb{C}$ by a projective toric variety X associated to the Newton polytope, P , of $D(z, \lambda)$ [15, Ch. 5]. The critical point equations become linear equations on X whose number of solutions is the degree of X . By Kushnirenko’s Theorem [19], this degree is the normalized volume of P , $(d+1)! \text{vol}(P)$. This bound is attained exactly when there are no solutions at infinity, which is the set $\partial X := X \setminus ((\mathbb{C}^\times)^d \times \mathbb{C})$ of points added in the compactification.

The compactified Bloch variety is a hypersurface in X . A *vertical face* of P is one that contains a segment parallel to the λ -axis. Corollary 3.9 shows that when P has no vertical faces, any solution on ∂X to the critical point equations is a singular point of the intersection of this hypersurface with ∂X . We state a simplified version of Corollary 3.9.

Theorem B. *If P has no vertical faces, then the bound of Corollary 2.5 is attained exactly when the compactified Bloch variety is smooth along ∂X .*

We give a class of graphs whose typical Bloch variety is smooth at infinity and whose Newton polytopes have no vertical faces. A periodic graph Γ is *dense* if it has every possible edge, given its support $\mathcal{A}(\Gamma)$ and fundamental domain W (see Section 4). The following is a consequence of Corollary 3.9 and Theorem 4.2.

Theorem C. *When $d = 2$ or 3 the Bloch variety of a generic operator on a dense periodic graph is smooth along ∂X , its Newton polytope has no vertical faces, and the bound of **Theorem A** is attained.*

Theorem C is an example of a recent trend in applications of algebraic geometry in which a highly structured optimization problem is shown to unexpectedly achieve a combinatorial bound on the number of critical points. A first instance was [9], which inspired [4] and [24].

Section 1 presents background on the spectrum of an operator on a periodic graph, and formulates our goal to bound the number of critical points of the function λ on the Bloch variety. At the beginning of Section 3, we recast extrema of the spectral band functions using the language of constrained optimization. Theorems A, B, and C are proven in Sections 2, 3, and 4. In Section 5, we use these results to prove the spectral edges conjecture for operators on three periodic graphs.

1. OPERATORS ON PERIODIC GRAPHS

Let d be a positive integer. We write $\mathbb{C}^\times := \mathbb{C} \setminus \{0\}$ for the multiplicative group of nonzero complex numbers and $\mathbb{T} := \{z \in \mathbb{C}^\times \mid |z| = 1\}$ for its maximal compact subgroup. Note that if $z \in \mathbb{T}$, then $\bar{z} = z^{-1}$. We write edges of a graph as pairs, (u, v) with u, v vertices, and understand that $(u, v) = (v, u)$.

1.1. Operators on periodic graphs. For more, see [2, Ch. 4]. A (\mathbb{Z}^d) -periodic graph is a simple (no multiple edges or loops) connected undirected graph Γ with a free cocompact action of \mathbb{Z}^d . Thus \mathbb{Z}^d acts freely on the vertices, $\mathcal{V}(\Gamma)$, and edges, $\mathcal{E}(\Gamma)$, of Γ preserving incidences, and \mathbb{Z}^d has finitely many orbits on each of $\mathcal{V}(\Gamma)$ and $\mathcal{E}(\Gamma)$. Figure 2 shows two




FIGURE 2. Two \mathbb{Z}^2 -periodic graphs.

\mathbb{Z}^2 -periodic graphs. One is the honeycomb lattice and the other is an abelian cover of K_4 , the complete graph on four vertices.

It is useful but not necessary to consider Γ immersed in \mathbb{R}^d so that \mathbb{Z}^d acts on Γ via translations. The graphs in Figure 2 are each immersed in \mathbb{R}^2 , and for each we show two independent vectors that generate the \mathbb{Z}^2 -action.

Choose a fundamental domain for this \mathbb{Z}^d -action whose boundary does not contain a vertex of Γ . In Figure 2, we have shaded the fundamental domains. Let W be the vertices of Γ lying in the fundamental domain. Then W is a set of representatives of \mathbb{Z}^d -orbits of $\mathcal{V}(\Gamma)$. Every \mathbb{Z}^d -orbit of edges contains one or two edges incident on vertices in W . An edge incident on W has the form $(u, a+v)$ for some $u, v \in W$ and $a \in \mathbb{Z}^d$. (If $a = 0$, then $u \neq v$

as Γ has no loops, and there are no restrictions when $a \neq 0$.) The support $\mathcal{A}(\Gamma)$ of Γ is the set of $a \in \mathbb{Z}^d$ such that $(u, a+v) \in \mathcal{E}(\Gamma)$ for some $u, v \in W$. This finite set depends on the choice of fundamental domain and it is centrally symmetric in that $\mathcal{A}(\Gamma) = -\mathcal{A}(\Gamma)$. As Γ is connected, the \mathbb{Z} -span of $\mathcal{A}(\Gamma)$ is \mathbb{Z}^d . For both graphs in Figure 2, this set consists of the columns of the matrix $\begin{pmatrix} 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & -1 \end{pmatrix}$.

A labeling of Γ is a pair of functions $e: \mathcal{E}(\Gamma) \rightarrow \mathbb{R}$ (edge weights) and $V: \mathcal{V}(\Gamma) \rightarrow \mathbb{R}$ (potential) that is \mathbb{Z}^d -invariant (constant on orbits). The set of labelings is the finite-dimensional vector space $\mathbb{R}^E \times \mathbb{R}^W$, where E is the set of orbits on $\mathcal{E}(\Gamma)$. Given a labeling $c = (e, V)$, we have the discrete operator L_c acting on functions f on $\mathcal{V}(\Gamma)$. Then $L_c(f)$ is defined by its value at $u \in \mathcal{V}(\Gamma)$,

$$L_c(f)(u) := V(u)f(u) + \sum_{(u,v) \in \mathcal{E}(\Gamma)} e_{(u,v)}(f(u) - f(v)).$$

We call L_c a *discrete periodic operator* on Γ , and may often omit the subscript c . It is a bounded self-adjoint operator on the Hilbert space $\ell^2(\Gamma)$ of square-summable functions on $\mathcal{V}(\Gamma)$, and has real spectrum.

1.2. Floquet theory. As the action of \mathbb{Z}^d on Γ commutes with the operator L , we may apply the Floquet transform, which reveals important structure of its spectrum. References for this *Floquet theory* include [2, 21, 22].

The Floquet (Fourier) transform is a linear isometry $\ell^2(\Gamma) \xrightarrow{\sim} L^2(\mathbb{T}^d, \mathbb{C}^W)$, from $\ell^2(\Gamma)$ to square-integrable functions on \mathbb{T}^d , the compact torus, with values in the vector space \mathbb{C}^W . The torus \mathbb{T}^d is the group of unitary characters of \mathbb{Z}^d . For $z \in \mathbb{T}^d$ and $a \in \mathbb{Z}^d$, the corresponding character value is the Laurent monomial

$$z^a := z_1^{a_1} z_2^{a_2} \cdots z_d^{a_d}.$$

The Floquet transform \hat{f} of a function f on $\mathcal{V}(\Gamma)$ is a function on $\mathbb{T}^d \times \mathcal{V}(\Gamma)$ such that for $z \in \mathbb{T}^d$ and $u \in \mathcal{V}(\Gamma)$,

$$(1) \quad \hat{f}(z, a+u) = z^a \hat{f}(z, u) \quad \text{for } a \in \mathbb{Z}^d.$$

Thus \hat{f} is determined by its values at the vertices W in the fundamental domain.

Let $\hat{f} \in L^2(\mathbb{T}^d, \mathbb{C}^W)$. Then for $u \in W$, $\hat{f}(u)$ is a function on \mathbb{T}^d . The action of the operator L on the Floquet transform \hat{f} is given by the formula

$$(2) \quad L(\hat{f})(u) = V(u)\hat{f}(u) + \sum_{(u,a+v) \in \mathcal{E}(\Gamma)} e_{(u,a+v)}(\hat{f}(u) - z^a \hat{f}(v)),$$

as $\hat{f}(a+v) = z^a \hat{f}(v)$. The exponents a which appear lie in the support $\mathcal{A}(\Gamma)$ of Γ . The simplicity of this expression is because L commutes with the \mathbb{Z}^d -action.

Thus in the standard basis for \mathbb{C}^W , the operator L becomes multiplication by a square matrix whose rows and columns are indexed by elements of W . Writing $\delta_{u,v}$ for the Kronecker delta function, the matrix entry in position (u, v) is the function

$$(3) \quad \delta_{u,v} \left(V(u) + \sum_{(u,w) \in \mathcal{E}(\Gamma)} e_{(u,w)} \right) - \sum_{(u,a+v) \in \mathcal{E}(\Gamma)} e_{(u,a+v)} z^a.$$

Example 1.1. Let Γ be the hexagonal lattice from Figure 2. Figure 3 shows a labeling in a neighborhood of its fundamental domain. Thus $W = \{u, v\}$ consists of two vertices and




FIGURE 3. A labeling of the hexagonal lattice.

there are three (orbits of) edges, with labels α, β, γ . Let $(x, y) \in \mathbb{T}^2$. The operator L is

$$\begin{aligned} L(\hat{f})(u) &= V(u)\hat{f}(u) + \alpha(\hat{f}(u) - \hat{f}(v)) + \beta(\hat{f}(u) - x^{-1}\hat{f}(v)) + \gamma(\hat{f}(u) - y^{-1}\hat{f}(v)), \\ L(\hat{f})(v) &= V(v)\hat{f}(v) + \alpha(\hat{f}(v) - \hat{f}(u)) + \beta(\hat{f}(v) - x\hat{f}(u)) + \gamma(\hat{f}(v) - y\hat{f}(u)). \end{aligned}$$

Collecting coefficients of $\hat{f}(u), \hat{f}(v)$, we represent L by the 2×2 -matrix,

$$(4) \quad L = \begin{pmatrix} V(u) + \alpha + \beta + \gamma & -\alpha - \beta x^{-1} - \gamma y^{-1} \\ -\alpha - \beta x - \gamma y & V(v) + \alpha + \beta + \gamma \end{pmatrix},$$

whose entries are Laurent polynomials in x, y . Notice that the support $\mathcal{A}(\Gamma)$ of Γ equals the set of exponents of monomials which appear in L . Observe that for $(x, y) \in \mathbb{T}^2$, $L^T = \overline{L}$, so that L is Hermitian, showing again that the operator L is self-adjoint. \diamond

What we saw in Example 1.1 holds in general. In the standard basis for \mathbb{C}^W , $L = L_c$ is multiplication by a $|W| \times |W|$ -matrix $L(z) = L_c(z)$ with each entry (3) a finite sum of monomials with exponents from $\mathcal{A}(\Gamma)$ (a *Laurent polynomial with support $\mathcal{A}(\Gamma)$*). Note that $(u, a+v) \in \mathcal{E}(\Gamma)$ if and only if $(-a+u, v) \in \mathcal{E}(\Gamma)$, these edges have the same label, and for $z \in \mathbb{T}^d$, $\overline{z^a} = z^{-a}$. Thus for $z \in \mathbb{T}^d$, the matrix is Hermitian, as $L(z)^T = L(z^{-1}) = \overline{L(z)}$.

1.3. Critical points of the Bloch variety. As $L(z)$ is Hermitian for $z \in \mathbb{T}^d$, its spectrum is real and consists of its $|W|$ eigenvalues

$$(5) \quad \lambda_1(z) \leq \lambda_2(z) \leq \cdots \leq \lambda_{|W|}(z).$$

These eigenvalues vary continuously with $z \in \mathbb{T}^d$, and $\lambda_j(z)$ is called the j th *spectral band function*, $\lambda_j: \mathbb{T}^d \rightarrow \mathbb{R}$. Its image is an interval in \mathbb{R} , called the j th *spectral band*. The eigenvalues (5) are the roots of the characteristic polynomial

$$(6) \quad D(z, \lambda) = D_c(z, \lambda) := \det(L_c(z) - \lambda I),$$

which we call the *dispersion function*. Its vanishing defines a hypersurface

$$(7) \quad \text{Var}(D_c(z, \lambda)) = \{(z, \lambda) \in \mathbb{T}^d \times \mathbb{R} \mid D(z, \lambda) = 0\},$$

called the *Bloch variety* of the operator L^1 . The Bloch variety is the union of $|W|$ branches with the j th branch equal to the graph of the j th spectral band function. The image of the Bloch variety under the projection to \mathbb{R} is the spectrum $\sigma(L)$ of the operator L . This projection is a function λ on the Bloch variety. Identifying the j th branch/graph with \mathbb{T}^d , the restriction of λ to that branch gives the corresponding spectral band function λ_j .

Figure 4 shows this for the operator L on the hexagonal lattice with edge weights 6, 3, 2 and zero potential V —for this we unfurl \mathbb{T}^2 , representing it by $[-\frac{\pi}{2}, \frac{3\pi}{2}]^2 \subset \mathbb{R}^2$, which is a fundamental domain in its universal cover. (That is, by quasimomenta in $[-\frac{\pi}{2}, \frac{3\pi}{2}]^2$.) It has




FIGURE 4. A Bloch variety and spectral bands for the hexagonal lattice.

two branches with each the graph of the corresponding spectral band function. An endpoint of a spectral band (*spectral edge*) is the image of an extremum of some band function $\lambda_j(z)$. For the hexagonal lattice at these parameters, each band function has two nondegenerate extrema, and these give the four spectral edges. These are also local extrema of the function λ on the Bloch variety.

The *spectral edges conjecture* [22, Conj. 5.25] for a periodic graph Γ asserts that for generic values of the parameters (e, V) , each spectral edge is attained by a single band, the extrema on the Bloch variety are isolated, and all extrema are nondegenerate (the spectral band function λ_j has a full rank Hessian matrix). Here, generic means that there is a nonconstant polynomial $p(e, V)$ in the parameters such that when $p(e, V) \neq 0$, these desired properties hold.

The entries in the matrix $L(z)$ and the function (6) defining the Bloch variety are all (Laurent) polynomials. In this setting it is natural to allow complex parameters, $e: \mathcal{E}(\Gamma) \rightarrow \mathbb{C}$, $V: \mathcal{V}(\Gamma) \rightarrow \mathbb{C}$ and variables $z \in (\mathbb{C}^\times)^d$, $\lambda \in \mathbb{C}$. With complex parameters and variables, $L_c(z)$ is no longer Hermitian, but it does satisfy $L_c(z)^T = L_c(z^{-1})$ and the Bloch variety is the complex algebraic hypersurface $\text{Var}(D_c(z, \lambda))$ in $(\mathbb{C}^\times)^d \times \mathbb{C}$ defined by the vanishing of the dispersion function $D_c(z, \lambda)$ of $L_c(z)$ (6).

In passing to the complex Bloch variety we may no longer distinguish branches $\lambda_j(z)$ of λ . At a smooth point (z_0, λ_0) whose projection z to $(\mathbb{C}^\times)^d$ is regular (in that $\frac{\partial D}{\partial \lambda}(z_0, \lambda_0) \neq 0$), there is a locally defined function f of z with $\lambda_0 = f(z_0)$ and $D(z, f(z)) = 0$ on its

¹This is also called the dispersion relation in the literature. We use the term Bloch variety as it is an algebraic variety and our perspective is to use methods from algebraic geometry in spectral theory.

domain, but this is not necessarily a global function of z . Consequently, we will consider the projection to the last coordinate to be a function λ on the Bloch variety, and then study its differential geometry, including its critical points.

Nondegeneracy of spectral edges is implied by the stronger condition that all critical points of the function λ on the complex Bloch variety are nondegenerate. Understanding the critical points of λ is a first step. Our aim is to bound the number of (isolated) critical points of λ on the Bloch variety of a given operator L , give criteria for when the bound is attained, prove that it is attained for generic operators on a class of graphs, and finally to use these results to prove the spectral edges conjecture for $2^{19} + 2$ graphs. We treat these in the following four sections.

2. BOUNDING THE NUMBER OF CRITICAL POINTS

We first recast extrema of spectral band functions in terms of constrained optimization. The complex Bloch variety is the hypersurface $\text{Var}(D(z, \lambda))$ in $(\mathbb{C}^\times)^d \times \mathbb{C}$ defined by the vanishing of the dispersion function $D(z, \lambda)$. **Critical points** of the function λ on the Bloch variety are points of the Bloch variety where the gradients in $(\mathbb{C}^\times)^d \times \mathbb{C}$ of λ and $D(z, \lambda)$ are linearly dependent. That is, a critical point is a point $(z, \lambda) \in (\mathbb{C}^\times)^d \times \mathbb{C}$ with $D(z, \lambda) = 0$ such that either the gradient $\nabla D(z, \lambda)$ vanishes or we have $\frac{\partial D}{\partial z_i}(z, \lambda) = 0$ for $i = 1, \dots, d$ and $\frac{\partial D}{\partial \lambda}(z, \lambda) \neq 0$ (as $\nabla \lambda = (0, \dots, 0, 1)$). In either case, we have

$$D(z, \lambda) = 0 \quad \text{and} \quad \frac{\partial D}{\partial z_i} = 0 \quad \text{for } i = 1, \dots, d.$$

Since $z_i \neq 0$, we obtain the equivalent system

$$(8) \quad D(z, \lambda) = z_1 \frac{\partial D}{\partial z_1} = \dots = z_d \frac{\partial D}{\partial z_d} = 0,$$

which we call the *critical point equations*.

Proposition 2.1. *A point $(z, \lambda) \in (\mathbb{C}^\times)^d \times \mathbb{C}$ is a critical point of the function λ on the Bloch variety $\text{Var}(D(z, \lambda))$ if and only if (8) holds.*

Proof. We already showed that at a critical point of λ , the equations (8) hold. Suppose now that $(z, \lambda) \in (\mathbb{C}^\times)^d \times \mathbb{C}$ is a solution to (8). As $D(z, \lambda) = 0$, the point lies on the Bloch variety. As $z \in (\mathbb{C}^\times)^d$, no coordinate z_i vanishes, which implies that $\frac{\partial D}{\partial z_i}(z, \lambda) = 0$ for $i = 1, \dots, d$. Thus the gradients $\nabla \lambda$ and ∇D are linearly dependent at (z, λ) , showing that it is a critical point. \square

Remark 2.2. A point $(z_0, \lambda_0) \in \mathbb{T}^d \times \mathbb{R}$ such that $\lambda_0 = \lambda_j(z_0)$ is an extreme value of the spectral band function λ_j is also a critical point of the Bloch variety. Indeed, either the gradient ∇D vanishes at (z_0, λ_0) or it does not vanish. If $\nabla D(z_0, \lambda_0) = \mathbf{0}$, then (z_0, λ_0) is a critical point. If $\nabla D(z_0, \lambda_0) \neq \mathbf{0}$, then the Bloch variety is smooth at (z_0, λ_0) and thus is a smooth point of the graph of λ_j . As $\lambda_0 = \lambda_j(z_0)$ is an extreme value of λ_j , the tangent plane is horizontal at (z_0, λ_0) . This implies that λ_j is differentiable (by the implicit function theorem) and that $\frac{\partial \lambda_j}{\partial z_i}(z_0, \lambda_0) = 0$ for $i = 1, \dots, d$. Thus the gradients of λ and D at (z_0, λ_0) are linearly dependent, showing that it is a critical point. \diamond

Bézout’s Theorem [29, Sect. 4.2.1] gives an upper bound on the number of isolated critical points: We may multiply each Laurent polynomial in (8) by a monomial to clear denominators and obtain ordinary polynomials. The product of their degrees is an upper bound for the number of the common zeroes that are isolated in the complex domain. Polyhedral bounds that exploit the structure of the Laurent polynomials are typically much smaller. Sources for these are [6, Ch. 7], [15, Ch. 5], and [30, Ch. 3]. These results bound the number of isolated common zeroes, counted with multiplicities. An isolated common zero z_0 of polynomials f_1, \dots, f_{d+1} on $(\mathbb{C}^\times)^d \times \mathbb{C}$ has multiplicity 1 exactly when the gradient of f_1, \dots, f_{d+1} spans the cotangent space at z_0 ; otherwise its multiplicity exceeds 1 (see [6, Ch. 4 Def. 2.1] and [7, Ch. 8.7 Def. 8]).

Let $\mathbb{C}[z^\pm, \lambda]$ be the ring of Laurent polynomials in z_1, \dots, z_d, λ where λ occurs with only nonnegative exponents. Note that $D(z, \lambda) \in \mathbb{C}[z^\pm, \lambda]$. The *support* $\mathcal{A}(\psi) \subset \mathbb{Z}^d \times \mathbb{N}$ of a polynomial $\psi \in \mathbb{C}[z^\pm, \lambda]$ is the set of exponents of monomials in ψ . The *Newton polytope* $\mathcal{N}(\psi) := \text{conv}(\mathcal{A}(\psi))$ of ψ is the convex hull of its support. Write $\text{vol}(\mathcal{N}(\psi))$ for the $(d+1)$ -dimensional Euclidean volume of the Newton polytope of ψ .

Example 2.3. We continue the example of the hexagonal lattice. Writing ℓ for $\alpha + \beta + \gamma - \lambda$, the dispersion function $D(x, y; \lambda)$ of the matrix (4) is

$$(V(u) + \ell)(V(v) + \ell) - (-\alpha - \beta x^{-1} - \gamma y^{-1})(-\alpha - \beta x - \gamma y).$$

In Figure 5 the monomials in $D(x, y; \lambda)$ label the columns of a 3×9 array which are their exponent vectors. Figure 5 also shows its Newton polytope, which has volume 2. \diamond




FIGURE 5. Support and Newton polytope of the hexagonal lattice operator.

Theorem 2.4. *For a polynomial $\psi \in \mathbb{C}[z^\pm, \lambda]$, the critical point equations for ψ*

$$(9) \quad \psi(z, \lambda) = z_1 \frac{\partial \psi}{\partial z_1} = \dots = z_d \frac{\partial \psi}{\partial z_d} = 0$$

have at most $(d+1)! \text{vol}(\mathcal{N}(\psi))$ isolated solutions in $(\mathbb{C}^\times)^d \times \mathbb{C}$, counted with multiplicity. When the bound is attained, all solutions are isolated.

We prove this at the end of the section.

As the Bloch variety is defined by the dispersion function $D(z, \lambda) = \det(L(z) - \lambda I)$, we deduce the following from Theorem 2.4.

Corollary 2.5. *The number of isolated critical points of the function λ on the Bloch variety for an operator L on a discrete periodic graph is at most $(d+1)! \text{vol}(\mathcal{N}(D))$.*

Theorem A follows from this and Lemma 4.1, which asserts that

$$\mathcal{N}(D) \subset |W|(\text{conv}(\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})),$$

where $\mathbf{e} = (0, \dots, 0, 1)$. This containment implies the inequality

$$(d+1)! \text{vol}(\mathcal{N}(D)) \leq (d+1)! |W|^{d+1} \text{vol}(\text{conv}(\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})) = d! |W|^{d+1} \text{vol}(\mathcal{A}(\Gamma)).$$

We prove Theorem 2.4 and Corollary 2.5 after developing some preliminary results.

2.1. A little algebraic geometry. For more from algebraic geometry, see [7, 29]. An (affine) *variety* is the set of common zeroes of some polynomials $f_1, \dots, f_r \in \mathbb{C}[x_1, \dots, x_n]$,

$$\text{Var}(f_1, \dots, f_r) := \{x \in \mathbb{C}^n \mid f_1(x) = \dots = f_r(x) = 0\}.$$

We also call this the set of *solutions* to the system $f_1 = \dots = f_r = 0$. We may replace any factor \mathbb{C} in \mathbb{C}^n by \mathbb{C}^\times , and then allow the corresponding variable to have negative exponents. The complement of a variety X is a (Zariski) open set. This defines the *Zariski topology* in which varieties are the closed sets. A variety is irreducible if it is not the union of two proper subvarieties. For an irreducible variety, any nonempty open set is dense (even in the classical topology) and any nonempty classically open set is dense in the Zariski topology. Maps $f: X \rightarrow Y \subset \mathbb{C}^m$ of varieties are given by m polynomials on X and the image $f(X)$ contains an open subset of its closure.

Suppose that $X = \text{Var}(f_1, \dots, f_r)$. The *smooth (nonsingular)* locus of X is the open subset of points of X where the Jacobian of f_1, \dots, f_r has maximal rank on X . Let f be a single polynomial. A point x is a smooth point on the hypersurface $\text{Var}(f)$ defined by f if $f(x) = 0$, so that $x \in \text{Var}(f)$ and if the gradient $\nabla f(x) = (\frac{\partial f}{\partial x_1}(x), \dots, \frac{\partial f}{\partial x_n}(x))$ is nonzero, so that some partial derivative of f does not vanish at x . The point $x \in \text{Var}(f)$ is *singular* if all partial derivatives of f vanish at x . The kernel of the Jacobian at $x \in X = \text{Var}(f_1, \dots, f_r)$ is the (Zariski) tangent space at x . The dimension of an irreducible variety is the dimension of a tangent space at any smooth point. An isolated point x of X has multiplicity one exactly when it is nonsingular.

Remark 2.6. Our definition of smooth and singular points of a variety depends upon its defining polynomials. For example, the variety defined by $(z - \lambda)^2$ is singular at every point. This *scheme-theoretic* notion of singularity is essential to our arguments in Sections 3 and 4, and is standard in algebraic geometry. \diamond

If X is irreducible, then any proper subvariety has smaller dimension. If $f: X \rightarrow Y$ is a map of varieties with $f(X)$ dense in Y , then there is an open subset U of Y such that if $y \in U$, then $\dim f^{-1}(y) + \dim Y = \dim X$. We also have *Bertini's Theorem*: if X is smooth, then U may be chosen so that for every $y \in U$, $f^{-1}(y)$ is smooth.

Projective space $\mathbb{P}(\mathbb{C}^n)$ is the set of one-dimensional linear subspaces (lines) of \mathbb{C}^n and is compact. It has dimension $n-1$ and subvarieties are given by homogeneous polynomials. The set U_0 of lines spanned by vectors whose initial coordinate is nonzero is isomorphic to \mathbb{C}^{n-1} under $v \mapsto \text{span}(1, v)$ and $\mathbb{P}(\mathbb{C}^n)$ is a compactification of $U_0 \simeq \mathbb{C}^{n-1}$.

2.2. Polyhedral bounds. The expression $(d+1)!\text{vol}(\mathcal{N}(\psi))$ of Theorem 2.4 is the *normalized volume* of $\mathcal{N}(\psi)$. This is Kushnirenko's bound [15, Ch. 6, Thm. 2.2] for the number of isolated solutions in $(\mathbb{C}^\times)^{d+1}$ to a system of $d+1$ polynomial equations, all with Newton polytope $\mathcal{N}(\psi)$. To prove Theorem 2.4, we first explain why Kushnirenko's bound applies to the system (9), and then why it bounds the number of isolated solutions on the larger space $(\mathbb{C}^\times)^d \times \mathbb{C}$.

For a monomial $z^a \lambda^j$ in $\mathbb{C}[z^\pm, \lambda]$, $a \in \mathbb{Z}^d$ and $j \in \mathbb{N}$. For each $i = 1, \dots, d$, this monomial is an eigenvector for the operator $z_i \frac{\partial}{\partial z_i}$ with eigenvalue a_i . Thus $\mathcal{A}(z_i \frac{\partial}{\partial z_i} \psi) \subset \mathcal{A}(\psi)$, giving the inclusion $\mathcal{N}(z_i \frac{\partial}{\partial z_i} \psi) \subset \mathcal{N}(\psi)$. A refined version of Kushnirenko's Theorem in which the polynomials may have different Newton polytopes is Bernstein's theorem [6, Sect. 7.5], which is in terms of a quantity called mixed volume, whose properties are developed in [10, Ch. IV]. The mixed volume of polytopes is monotone under inclusion of polytopes and it equals the normalized volume when all polytopes coincide. It follows that the theorems of Bernstein and Kushnirenko together give the bound of $(d+1)!\text{vol}(\mathcal{N}(\psi))$ for the number of isolated solutions to the system (9) in $(\mathbb{C}^\times)^{d+1}$. To extend this to solutions in the larger space $(\mathbb{C}^\times)^d \times \mathbb{C}$, we develop some theory of projective toric varieties.

2.3. Projective toric varieties. For Kushnirenko's Theorem and our extension, we replace the nonlinear equations (9) on $(\mathbb{C}^\times)^d \times \mathbb{C}$ by linear equations on a projective variety. We follow the discussion of [30, Ch. 3]. Let $f \in \mathbb{C}[z^\pm, \lambda]$ be a polynomial with support $\mathcal{A} = \mathcal{A}(f)$. To simplify the presentation, we will at times assume that the origin $\mathbf{0}$ lies in \mathcal{A} . The results hold without this assumption, as explained in [30, Ch. 3].

Writing $\mathbb{C}^{\mathcal{A}}$ for the vector space with basis indexed by elements of \mathcal{A} , consider the map

$$\begin{aligned} \varphi_{\mathcal{A}} : (\mathbb{C}^\times)^d \times \mathbb{C} &\longrightarrow \mathbb{C}^{\mathcal{A}} \\ (z, \lambda) &\longmapsto (z^a \lambda^j \mid (a, j) \in \mathcal{A}) . \end{aligned}$$

This map linearizes nonlinear polynomials. Indeed, write f as a sum of monomials,

$$f = \sum_{(a,j) \in \mathcal{A}} c_{(a,j)} z^a \lambda^j .$$

If $\{x_{(a,j)} \mid (a, j) \in \mathcal{A}\}$ are variables (coordinate functions) on $\mathbb{C}^{\mathcal{A}}$, then

$$(10) \quad \Lambda_f := \sum_{(a,j) \in \mathcal{A}} c_{(a,j)} x_{(a,j)}$$

is a linear form on $\mathbb{C}^{\mathcal{A}}$, and we have $f(z, \lambda) = \Lambda_f(\varphi_{\mathcal{A}}(z, \lambda)) =: \varphi_{\mathcal{A}}^*(\Lambda_f)$.

Since $\mathbf{0} \in \mathcal{A}$, the corresponding coordinate $x_{\mathbf{0}}$ of $\varphi_{\mathcal{A}}$ is 1 and so the image of $\varphi_{\mathcal{A}}$ lies in the principal affine open subset $U_{\mathbf{0}}$ of the projective space $\mathbb{P}^{\mathcal{A}} := \mathbb{P}(\mathbb{C}^{\mathcal{A}}) = \mathbb{P}^{|\mathcal{A}|-1}$. This is the subset of $\mathbb{P}^{\mathcal{A}}$ where $x_{\mathbf{0}} \neq 0$ and it is isomorphic to the affine space $\mathbb{C}^{|\mathcal{A}|-1}$. We define $X_{\mathcal{A}}$ to be the closure of the image $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^{d+1})$ in the projective space $\mathbb{P}^{\mathcal{A}}$, which is a projective toric variety. Because the map $\varphi_{\mathcal{A}}$ is continuous on $(\mathbb{C}^\times)^d \times \mathbb{C}$, $X_{\mathcal{A}}$ is also the closure of the image $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C})$.

The map $\varphi_{\mathcal{A}}$ is not necessarily injective; we describe its fibers. Let $\mathbb{Z}\mathcal{A} \subset \mathbb{Z}^{d+1}$ be sublattice generated by all differences $\alpha - \beta$ for $\alpha, \beta \in \mathcal{A}$. When $\mathbf{0} \in \mathcal{A}$ this is the sublattice generated by \mathcal{A} , and it has full rank $d + 1$ if and only if $\text{conv}(\mathcal{A})$ has full dimension $d + 1$.

Let $G_{\mathcal{A}}$ be $\text{Hom}(\mathbb{Z}^{d+1}/\mathbb{Z}\mathcal{A}, \mathbb{C}^\times) \subset (\mathbb{C}^\times)^{d+1}$, which acts on $(\mathbb{C}^\times)^d \times \mathbb{C}$. The fibers of $\varphi_{\mathcal{A}}$ are exactly the orbits of $G_{\mathcal{A}}$ on $(\mathbb{C}^\times)^d \times \mathbb{C}$. If $\text{conv}(\mathcal{A})$ does not have full dimension, then $G_{\mathcal{A}}$ has positive dimension as do all fibers of $\varphi_{\mathcal{A}}$, otherwise $G_{\mathcal{A}}$ is a finite group and $\varphi_{\mathcal{A}}$ has finite fibers. On the torus $(\mathbb{C}^\times)^{d+1}$, $G_{\mathcal{A}}$ acts freely and $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^{d+1})$ is identified with $(\mathbb{C}^\times)^{d+1}/G_{\mathcal{A}}$. To describe the fibers of $\varphi_{\mathcal{A}}$ on $(\mathbb{C}^\times)^d \times \{0\} = ((\mathbb{C}^\times)^d \times \mathbb{C}) \setminus (\mathbb{C}^\times)^{d+1}$, note that $(\mathbb{C}^\times)^{d+1}$ acts on this through the homomorphism π that sends its last (λ) coordinate to $\{1\}$. Thus the fibers of $\varphi_{\mathcal{A}}$ on $(\mathbb{C}^\times)^d \times \{0\}$ are exactly the orbits of $\pi(G_{\mathcal{A}}) \subset (\mathbb{C}^\times)^d$.

Proposition 2.7. *The dimension of $X_{\mathcal{A}}$ is the dimension of $\text{conv}(\mathcal{A})$. The fibers of $\varphi_{\mathcal{A}}$ on $(\mathbb{C}^\times)^{d+1}$ are the orbits of $G_{\mathcal{A}}$ and its fibers on $(\mathbb{C}^\times)^d \times \{0\}$ are the orbits of $\pi(G_{\mathcal{A}})$.*

We return to the situation of Theorem 2.4. Let $\psi \in \mathbb{C}[z^\pm, \lambda]$ be a polynomial with support \mathcal{A} . As each polynomial in (9) has support a subset of \mathcal{A} , each corresponds to a linear form on $\mathbb{P}^{\mathcal{A}}$ as in (10). The corresponding system of linear forms defines a linear subspace M_ψ of $\mathbb{P}^{\mathcal{A}}$. We have the following proposition (a version of [30, Lemma 3.5]).

Proposition 2.8. *The solutions to (9) are the inverse images under $\varphi_{\mathcal{A}}$ of points in the linear section $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C}) \cap M_\psi$. When $\varphi_{\mathcal{A}}$ is an injection, it is a bijection between solutions to (9) on $(\mathbb{C}^\times)^d \times \mathbb{C}$ and points in $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C}) \cap M_\psi$.*

Proof of Theorem 2.4. When $\text{vol}(\mathcal{N}(\psi)) = 0$, so that $\mathcal{N}(\psi)$ does not have full dimension $d + 1$, then each fiber of $\varphi_{\mathcal{A}}$ is positive-dimensional and so by Proposition 2.8 there are no isolated solutions to (9).

Suppose that $\text{vol}(\mathcal{N}(\psi)) > 0$. Then every fiber of $\varphi_{\mathcal{A}}$ is an orbit of the finite group $G_{\mathcal{A}}$. Over points of $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^{d+1})$, each fiber consists of $|G_{\mathcal{A}}|$ points and over $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \{0\})$ each fiber consists of $|\pi(G_{\mathcal{A}})| \leq |G_{\mathcal{A}}|$ points. As $X_{\mathcal{A}}$ is the closure of $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C})$, the number of isolated points in $X_{\mathcal{A}} \cap M_\psi$ is at least the number of isolated points in $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C}) \cap M_\psi$, both counted with multiplicity. The degree of the projective variety $X_{\mathcal{A}}$ is an upper bound for the number of isolated points in $X_{\mathcal{A}} \cap M_\psi$, which is explained in [30, Ch. 3.3]. There, the product of $|G_{\mathcal{A}}|$ and the degree of $X_{\mathcal{A}}$ is shown to be $(d+1)!\text{vol}(\mathcal{N}(\psi))$, the normalized volume of the Newton polytope of ψ . This gives the bound of Theorem 2.4. That all points are isolated when the bound of the degree is attained is Proposition 3.2 in the next section. \square

3. PROOF OF THEOREM B

We give conditions for when the upper bound of Corollary 2.5 is attained. By Proposition 2.1, the critical points of the function λ on the Bloch variety $\text{Var}(D)$ are the solutions in $(\mathbb{C}^\times)^d \times \mathbb{C}$ to the critical point equations (8). Let $\mathcal{A} = \mathcal{A}(D)$ be the support of the polynomial D . The critical points are $\varphi_{\mathcal{A}}^{-1}(X_{\mathcal{A}} \cap M_D)$, where $X_{\mathcal{A}} \subset \mathbb{P}^{\mathcal{A}}$ is the closure of $\varphi_{\mathcal{A}}((\mathbb{C}^\times)^d \times \mathbb{C})$ and M_D is the subspace of $\mathbb{P}^{\mathcal{A}}$ defined by linear forms corresponding (as in (10)) to the polynomials in (8). For the bound of Theorem 2.4 and Corollary 2.5, note that the number of isolated points of $X_{\mathcal{A}} \cap M_D$ is at most the product of the degree of $X_{\mathcal{A}}$ with the cardinality of a fiber of $\varphi_{\mathcal{A}}$, which is $(d+1)!\text{vol}(\mathcal{N}(D))$. We establish Theorem B concerning the sharpness of this bound by characterizing when the inequality of Theorem 2.4 is strict and then interpreting that for the critical point equations.

Remark 3.1. Let $X \subset \mathbb{P}^n$ be a variety of dimension d and $M \subset \mathbb{P}^n$ a linear subspace of codimension d . The number of points in $X \cap M$ does not depend on M when the intersection is transverse; it is the *degree* of X [29, p. 234]. When the intersection is not transverse, intersection theory gives a refinement [14, Ch. 6]. For each irreducible component Z of the intersection $X \cap M$, there is a positive integer—the intersection multiplicity along Z —such that the sum of these multiplicities is the degree of X . When Z is positive-dimensional this number is the degree of a zero-cycle constructed on Z (it is at least the degree of Z) and when Z is zero-dimensional (a point), it is the local multiplicity [29, Ch. 4]. \diamond

A consequence of Remark 3.1 is the following.

Proposition 3.2. *Let X, M be as in Remark 3.1. The number (counted with multiplicity) of isolated points of $X \cap M$ is strictly less than the degree of X if and only if the intersection has a positive-dimensional component.*

Write $X_{\mathcal{A}}^{\circ} := \varphi_{\mathcal{A}}((\mathbb{C}^{\times})^d \times \mathbb{C})$ for the image of $\varphi_{\mathcal{A}}$ and $\partial X_{\mathcal{A}} := X_{\mathcal{A}} \setminus X_{\mathcal{A}}^{\circ}$, the points of $X_{\mathcal{A}}$ added to $X_{\mathcal{A}}^{\circ}$ when taking the closure. This is the *boundary* of $X_{\mathcal{A}}$. In the Introduction, points of $\partial X_{\mathcal{A}}$ were referred to as ‘lying at infinity’.

Corollary 3.3. *For a polynomial $\psi \in \mathbb{C}[z^{\pm}, \lambda]$, the inequality of Theorem 2.4 is strict if and only if $\partial X_{\mathcal{A}} \cap M_{\psi} \neq \emptyset$.*

Proof. The inequality of Theorem 2.4 is strict if either of the following hold.

- (1) $X_{\mathcal{A}} \cap M_{\psi}$ has an isolated point not lying in $X_{\mathcal{A}}^{\circ}$.
- (2) $X_{\mathcal{A}} \cap M_{\psi}$ contains a positive-dimensional component Z .

In (1), $X_{\mathcal{A}} \cap M_{\psi}$ has isolated points in $\partial X_{\mathcal{A}} \cap M_{\psi}$, so the intersection is nonempty. In (2), Z is a projective variety of dimension at least one. The set $X_{\mathcal{A}}^{\circ}$ is an affine variety, and we cannot have $Z \subset X_{\mathcal{A}}^{\circ}$ as the only projective varieties that are also subvarieties of an affine variety are points. Thus $Z \cap \partial X_{\mathcal{A}} \neq \emptyset$, which completes the proof. \square

3.1. Facial systems. We return to the general case of a toric variety. Let $\mathcal{A} \subset \mathbb{Z}^n$ be a finite set of points with corresponding projective toric variety $X_{\mathcal{A}} \subset \mathbb{P}^A$. We have the following description of the points of its boundary, $X_{\mathcal{A}} \setminus \varphi_{\mathcal{A}}((\mathbb{C}^{\times})^n)$.

Let $P := \text{conv}(\mathcal{A})$, the convex hull of \mathcal{A} . The dot product with a nonzero vector $w \in \mathbb{R}^n$, $a \mapsto w \cdot a$, defines a linear function on \mathbb{R}^n . For $w \in \mathbb{R}^n$, set $h(w) := \min\{w \cdot a \mid a \in P\}$. The set $F = \{p \in P \mid w \cdot p = h(w)\}$ of minimizers is the *face* of P *exposed* by w . We have that $F = \text{conv}(F \cap \mathcal{A})$, and may write \mathcal{F} for $F \cap \mathcal{A}$. As $\mathcal{A} \subset \mathbb{Z}^n$, we only need integer vectors $w \in \mathbb{Z}^n$ to expose all faces of P . If $\dim F = \dim P - 1$, then F is a *facet*.

For each face F of P , there is a corresponding coordinate subspace $\mathbb{P}^{\mathcal{F}}$ of \mathbb{P}^A —this is the set of points $z = [z_a \mid a \in \mathcal{A}] \in \mathbb{P}^A$ such that $a \notin F$ implies that $z_a = 0$. The image of the map $\varphi_{\mathcal{F}}: (\mathbb{C}^{\times})^n \rightarrow \mathbb{P}^{\mathcal{F}} \subset \mathbb{P}^A$ has closure the toric variety $X_{\mathcal{F}}$. Its dimension is equal to the dimension of the face F . Write $X_{\mathcal{F}}^{\circ}$ for the image of $\varphi_{\mathcal{F}}$. This description and the following proposition is essentially [15, Prop. 5.1.9].

Proposition 3.4. *The boundary of the toric variety $X_{\mathcal{A}}$ is the disjoint union of the sets $X_{\mathcal{F}}^{\circ}$ for all the proper faces F of $\text{conv}(\mathcal{A})$.*

Let $f = \sum_{a \in \mathcal{A}} c_a x^a$ be a polynomial with support \mathcal{A} . We observed that if Λ is the corresponding linear form (10) on \mathbb{P}^A , then the variety $\text{Var}(f) \subset (\mathbb{C}^\times)^n$ of f is the pullback along $\varphi_{\mathcal{A}}$ of $X_{\mathcal{A}}^\circ \cap M$, where $M := \text{Var}(\Lambda)$ is the hyperplane defined by Λ . Let F be a proper face of P . Then $X_{\mathcal{F}}^\circ \cap M$ pulls back along $\varphi_{\mathcal{F}}$ to the variety of

$$\varphi_{\mathcal{F}}^{-1}(\Lambda) = \sum_{a \in F} c_a x^a$$

in $(\mathbb{C}^\times)^n$. This sum of the terms of f whose exponents lie in F is a *facial form* of f and is written $f|_F$. Given a system $\Phi: f_1 = \cdots = f_n = 0$ involving Laurent polynomials with support \mathcal{A} , the system $f_1|_F = \cdots = f_n|_F = 0$ of their facial forms is the *facial system* $\Phi|_F$ of Φ .

Corollary 3.5. *Let M be the intersection of the hyperplanes given by the polynomials in a system Φ of Laurent polynomials with support \mathcal{A} . For each face F of $\text{conv}(\mathcal{A})$, the points of $X_{\mathcal{F}}^\circ \cap M$ pull back under $\varphi_{\mathcal{F}}$ to the solutions of the facial system $\Phi|_F$.*

If no facial system $\Phi|_F$ has a solution, then the number of solutions to $\Phi = 0$ on $(\mathbb{C}^\times)^n$ is $n! \text{vol}(\text{conv}(\mathcal{A}))$.

Proof. The first statement follows from the observation about a single polynomial f and its facial form $f|_F$, and the second is a consequence of a version of Corollary 3.3 for $X_{\mathcal{A}} \setminus \varphi_{\mathcal{A}}((\mathbb{C}^\times)^n)$. \square

The second statement is essentially [3, Thm. B] and is also explained in [30, Sect. 3.4].

3.2. Facial systems of the critical point equations. We prove Theorem B from the Introduction by interpreting the facial systems of the critical point equations. It is useful to introduce the following notion. A polynomial $f(x)$ in $x \in (\mathbb{C}^\times)^n$ is *quasi-homogeneous* with *quasi-homogeneity* $w \in \mathbb{Z}^n$ if there is a number $0 \neq w_f$ such that

$$a \in \mathcal{A}(f) \implies w \cdot a = w_f.$$

Equivalently, f is quasi-homogeneous if its support $\mathcal{A}(f)$ lies on a hyperplane not containing the origin. The quasi-homogeneities of f are those $w \in \mathbb{Z}^n$ whose dot product is constant on $\mathcal{A}(f)$. For $t \in \mathbb{C}^\times$ and $w \in \mathbb{Z}^n$, let $t^w := (t^{w_1}, \dots, t^{w_n}) \in (\mathbb{C}^\times)^n$.

Lemma 3.6. *Suppose that f has a quasi-homogeneity $w \in \mathbb{Z}^n$. Then*

- (1) *For $t \in \mathbb{C}^\times$ and $x \in (\mathbb{C}^\times)^n$, we have $f(t^w \cdot x) = t^{w_f} f(x)$.*
- (2) *We have*

$$w_f f = \sum_{i=1}^n w_i x_i \frac{\partial f}{\partial x_i}.$$

Proof. Note that for $a \in \mathbb{Z}^n$, $(t^w \cdot x)^a = t^{w \cdot a} x^a$. The first statement follows. For the second, note that $a_i x^a = x_i \frac{\partial}{\partial x_i} x^a$. \square

Let $\psi \in \mathbb{C}[z^\pm, \lambda]$ have support $\mathcal{A} \subset \mathbb{Z}^d \times \mathbb{N}$ and Newton polytope $P := \text{conv}(\mathcal{A})$. We will assume that P has dimension $d+1$, and also that $\mathcal{A} \cap \mathbb{Z}^d \times \{0\}$ is a facet of \mathcal{A} , called its *base*. Let (9) be the critical point equations for λ on ψ and $M_\psi \subset \mathbb{P}^A$ the corresponding linear subspace of codimension $d+1$.

Let $\mathbf{0} := 0^d$ in \mathbb{Z}^d and $\mathbf{e} := (\mathbf{0}, 1)$. The base of \mathcal{A} is exposed by \mathbf{e} and it is the support of $\psi(z, 0)$. A main difference between the sparse equations of Section 3.1 and the critical point equations (8) is that the critical point equations allow solutions with $\lambda = 0$, which is the component of the boundary of the toric variety corresponding to the base of \mathcal{A} . A face F of P is *vertical* if it contains a vertical line segment, one parallel to \mathbf{e} .

Lemma 3.7. *Suppose that F is a proper face of P that is not the base of P and is not vertical. Then the corresponding facial system of the critical point equations has a solution if and only if the hypersurface $\text{Var}(\psi|_F)$ defined by $\psi|_F$ in $(\mathbb{C}^\times)^{d+1}$ is singular.*

Proof. Let $0 \neq w \in \mathbb{Z}^{d+1}$ be an integer vector that exposes the face F . As F is not vertical we may assume that w_{d+1} is nonzero. As F is not the base, it lies on an affine hyperplane that does not contain the origin, so that $\psi|_F$ is quasi-homogeneous with some quasi-homogeneity w . Write w_F for the constant $w \cdot a$ for $a \in F$. By Lemma 3.6 (2), we have

$$(11) \quad w_F \psi|_F = \sum_{i=1}^d w_i z_i \frac{\partial \psi|_F}{\partial z_i} + w_{d+1} \lambda \frac{\partial \psi|_F}{\partial \lambda}.$$

Suppose now that (z, λ) is a solution of the restriction of the critical point equations to the face F . That is, at (z, λ) ,

$$\psi|_{\mathcal{F}} = \left(z_1 \frac{\partial \psi}{\partial z_1} \right) \Big|_F = \cdots = \left(z_d \frac{\partial \psi}{\partial z_d} \right) \Big|_F = 0.$$

Observe that $(z_i \frac{\partial \psi}{\partial z_i})|_F = z_i \frac{\partial \psi|_F}{\partial z_i}$ (and the same for λ). Since $w_{d+1} \neq 0$, these equations and (11) together imply that $(\lambda \frac{\partial \psi}{\partial \lambda})|_F = 0$, which implies that (z, λ) is a singular point of the hypersurface $\text{Var}(\psi|_F)$ defined by $\psi|_F$. \square


We deduce the following theorem.

Theorem 3.8. *If the Newton polytope $\mathcal{N}(\psi)$ of ψ has no vertical faces and the restriction of ψ to each face that is not the base of $\mathcal{N}(\psi)$ defines a smooth variety, then the critical point equations have exactly $(d+1)!\text{vol}(\mathcal{N}(\mathcal{A}))$ solutions in $(\mathbb{C}^\times)^d \times \mathbb{C}$.*

We apply this when ψ is the dispersion function $D(z, \lambda)$. Recall that the boundary of the variety $X_{\mathcal{A}}$ (X_D) corresponds to all proper faces of its Newton polytope $\mathcal{N}(D)$, except for its base. We deduce the following precise version of **Theorem B**.

Corollary 3.9. *Let L be an operator on a periodic graph and set $D = \det(L(z) - \lambda I)$. If $\mathcal{N}(D)$ has no vertical faces and if for each face F that is not its base, $\text{Var}(D|_F)$ is smooth, then the Bloch variety has exactly $(d+1)!\text{vol}(\text{conv}(\mathcal{A}(D)))$ critical points.*

Example 3.10. The restriction on vertical faces is necessary. General operators on the second graph in Figure 2 (an abelian cover of K_4) have the following Newton polytope:



It has base $[-1, 1]^2$, apex $(0, 0, 4)$, and the remaining vertices are at $(\pm 1, 0, 1)$ and $(0, \pm 1, 1)$. It has volume $20/3$, so we expect $40 = 3! \cdot 20/3$ critical points. However, there are at most 32 critical points, as direct computation shows that the critical point equations have two solutions on each of its four vertical faces. \diamond

4. NEWTON POLYTOPES AND DENSE PERIODIC GRAPHS

The Newton polytope $\mathcal{N}(D)$ of the dispersion function of an operator on a periodic graph is central to our results. In Section 4.1 we associate a polytope $\mathcal{N}(\Gamma)$ to any periodic graph Γ such that $\mathcal{N}(D) \subset \mathcal{N}(\Gamma)$ for any operator on Γ , and that we have equality for almost all parameter values. We call $\mathcal{N}(\Gamma)$ the *Newton polytope* of Γ .

A periodic graph Γ is dense if it has every possible edge, given its support $\mathcal{A}(\Gamma)$ and fundamental domain W . Every periodic graph is a subgraph of a minimal dense periodic graph. We identify the Newton polytope of a dense periodic graph and show that when $d = 2$ or 3 , a general operator on Γ satisfies Corollary 3.9, which implies Theorem C.

Let Γ be a connected \mathbb{Z}^d -periodic graph with fundamental domain W . Its support $\mathcal{A}(\Gamma)$ is the finite set of points $a \in \mathbb{Z}^d$ such that there is an edge between W and $a+W$. The integer span of $\mathcal{A}(\Gamma)$ is \mathbb{Z}^d , as Γ is connected. The graph Γ is *dense* if for every $a \in \mathcal{A}(\Gamma)$, there is an edge in Γ between every pair of vertices in the union of W and $a+W$. In particular, the restriction of Γ to W is the complete graph on W . The graphs of Figures 1 and 6 are dense, while those of Figure 2 are not dense.




FIGURE 6. A dense graph Γ and its support $\mathcal{A}(\Gamma)$ with convex hull.




FIGURE 7. Newton polytopes of dense graphs.

pyramid with base $m \operatorname{conv}(\mathcal{A}(\Gamma))$ and apex me , and it has no vertical faces.

Theorem 4.2. *Let Γ be a dense \mathbb{Z}^d -periodic graph. There is a nonempty Zariski open subset U of the parameter space Y such that for $c \in U$, the Newton polytope of $D_c(z, \lambda)$ is the pyramid mQ . When $d = 2$ or 3 , then we may choose U so that for every $c \in U$ and face F of mQ that is not its base, $\operatorname{Var}(D_c|_F)$ is smooth.*

Together with Corollary 3.9, this implies Theorem C from the Introduction. We prove Theorem 4.2 in the following two subsections.

4.1. The Newton polytope of Γ . For a periodic graph Γ , the space of parameters (e, V) for operators on Γ is $Y = \mathbb{C}^E \times \mathbb{C}^W$. Treating parameters as indeterminates gives the *generic dispersion function* $D(e, V, z, \lambda)$, which is a polynomial in z, λ whose coefficients are polynomials in the parameters e, V . The *Newton polytope* $\mathcal{N}(\Gamma)$ of Γ is the convex hull of the monomials in z, λ that appear in $D(e, V, z, \lambda)$.

Lemma 4.3. *For $c \in Y$, $\mathcal{N}(D_c(z, \lambda))$ is a subpolytope of $\mathcal{N}(\Gamma)$. The set of $c \in Y$ such that $\mathcal{N}(D_c(z, \lambda)) = \mathcal{N}(\Gamma)$ is a dense open subset U . When Γ is a dense periodic graph, $\mathcal{N}(\Gamma) = mQ$.*

Proof. For any $c = (e, V) \in Y$, $D_c(z, \lambda)$ is the evaluation of the generic dispersion function $D(e, V, z, \lambda)$ at the point (e, V) . Thus $\mathcal{N}(D_c) \subset \mathcal{N}(\Gamma)$.

The coefficient $C_{(a,j)}$ of a monomial $z^a \lambda^j$ in $D(e, V, z, \lambda)$ is a polynomial in (e, V) . For any $c = (e, V) \in Y$, $z^a \lambda^j$ appears in D_c if and only if $C_{(a,j)}(e, V) \neq 0$. Thus, we have the

equality $\mathcal{N}(D_c) = \mathcal{N}(\Gamma)$ of Newton polytopes if and only if $C_{(a,j)}(e, V) \neq 0$ for every vertex (a, j) of $\mathcal{N}(\Gamma)$, which defines a dense open subset $U \subset Y$.

When Γ is dense and no parameter c vanishes, then every diagonal entry of $L_c(z) - \lambda I$ has support $\mathcal{A}(\Gamma) \cup \{\mathbf{e}\}$. This implies that $\mathcal{N}(\Gamma) = mQ$. \square

4.2. Smoothness of the Bloch variety at infinity. Let Γ be a dense periodic graph with $d = 2$ or 3 . Let $U \subset Y$ be the subset of Lemma 4.3. We show that for each face F of $\mathcal{N}(\Gamma)$ that is not its base, there is a nonempty open subset U_F of U such that for $c \in U_F$, the restriction $D_c|_F$ to the monomials in F defines a smooth hypersurface. Then for parameters c in the intersection of the U_F , the operator satisfies the hypotheses of Corollary 3.9, which proves Theorem 4.2 and Theorem C.

Let F be a face of $\mathcal{N}(\Gamma)$ that is not its base and let $c \in U$. We may assume that F is not a vertex, for then $D_c|_F$ is a single term and $\text{Var}(D_c|_F) = \emptyset$. Since $\mathcal{N}(\Gamma) = mQ$, there is a unique face G of Q such that $F = mG$. We have that

$$D_c(z, \lambda)|_F = \det((L_c(z) - \lambda I)|_G),$$

where each entry of the matrix $(L_c(z) - \lambda I)|_G$ is the facial form $f|_G$ of the corresponding entry f of $L_c(z) - \lambda I$.

Since G is not the base of Q (and thus does not contain the origin), we make the following observation, which follows from the form of the operator L_c (2). If the apex $\mathbf{e} = (\mathbf{0}, 1)$ of Q lies in G and f is a diagonal entry of $(L_c(z) - \lambda I)|_G$, then f contains the term $-\lambda$. Any other integer point $a \in G$ is nonzero and lies in the support $\mathcal{A}(\Gamma)$ of Γ , and the coefficient of z^a in f is $-e_{(u, a+v)}$, where f is the entry in row u and column v . Consequently, except possibly for terms $-\lambda$, all coefficients of entries in $(L_c(z) - \lambda I)|_G$ are distinct parameters.

Suppose that the fundamental domain is $W = \{v_1, \dots, v_m\}$ so that we may index the rows and columns of $L_c(z)$ by $1, \dots, m$. Let $Y' \subset Y$ be the set of parameters c where

$$e_{(v_i, a+v_j)} = 0 \quad \text{if } a \in G \text{ and } j \neq i, i+1.$$

(Here, $m+1$ is interpreted to be 1.) For $c \in Y'$, all entries of $L_c(z)|_G$ are zero, except on the diagonal, the first super diagonal, and the lower left entry. The same arguments as in the proof of Lemma 4.3 show that there exist parameters $c \in Y'$ such that $D_c(z, \lambda)$ has Newton polytope $\mathcal{N}(\Gamma)$. Thus $Y' \cap U \neq \emptyset$, where $U \subset Y$ is the set of Lemma 4.3.

Theorem 4.4. *There exists an open subset U' of Y' with $U' \subset U$ such that if $c \in U'$, then $\text{Var}(D_c(z, \lambda)|_F)$ is a smooth hypersurface in $(\mathbb{C}^\times)^{d+1}$.*

Since smoothness of $\text{Var}(D_c(z, \lambda)|_F)$ is an open condition on the space Y of parameters, this will complete the proof of Theorem 4.2, and thus also of Theorem C.

Proof. Let us write $\psi_c(z, \lambda)$ for the facial polynomial $D_c(z, \lambda)|_F$. We will show that the set of $c \in Y'$ such that $\text{Var}(\psi_c(z, \lambda))$ is singular is a finite union of proper algebraic subvarieties. As $c \in Y'$, the only nonzero entries in the matrix $(L_c(z) - \lambda I)|_G$ are its diagonal entries $f_1(z, \lambda), \dots, f_m(z, \lambda)$ and the entries $g_1(z), \dots, g_m(z)$ which are in positions $(1, 2), \dots, (m-1, m)$ and $(m, 1)$, respectively. Thus

$$\psi_c(z, \lambda) = D_c(z, \lambda)|_F = \det((L_c(z) - \lambda I)|_G) = \prod_{i=1}^m f_i(z, \lambda) - (-1)^m \prod_{i=1}^m g_i(z).$$

For a polynomial f in the variables (z, λ) , write $\nabla_{\mathbb{T}}$ for the toric gradient operator,

$$\nabla_{\mathbb{T}}f := \left(z_1 \frac{\partial f}{\partial z_1}, \dots, z_d \frac{\partial f}{\partial z_d}, \lambda \frac{\partial f}{\partial \lambda} \right).$$

Note that

$$(12) \quad \nabla_{\mathbb{T}}\psi_c = \sum_{i=1}^m (\nabla_{\mathbb{T}}f_i) f_1 \cdots \widehat{f}_i \cdots f_m - (-1)^m \sum_{i=1}^m (\nabla_{\mathbb{T}}g_i) g_1 \cdots \widehat{g}_i \cdots g_m.$$

Here \widehat{f}_i indicates that f_i does not appear in the product, and the same for \widehat{g}_i .

Let $(z, \lambda) \in \text{Var}(\psi_c)$ be a singular point. Then $\psi_c(z, \lambda) = 0$ and $\nabla_{\mathbb{T}}\psi_c(z, \lambda) = \mathbf{0}$. There are five cases that depend upon the number of polynomials f_i, g_j vanishing at (z, λ) .

- (i) At least two polynomials f_p and f_q and two polynomials g_r and g_s vanish at (z, λ) . Thus $\psi(z, \lambda) = 0$ and by (12) this implies that $\nabla_{\mathbb{T}}\psi_c(z, \lambda) = \mathbf{0}$.
- (ii) At least two polynomials f_p and f_q and exactly one polynomial g_s vanish at (z, λ) . Thus $\psi(z, \lambda) = 0$ and by (12) if $\nabla_{\mathbb{T}}\psi_c(z, \lambda) = \mathbf{0}$, then $\nabla_{\mathbb{T}}g_s(z, \lambda) = \mathbf{0}$.
- (iii) Exactly one polynomial f_p and at least two polynomials g_r and g_s vanish at (z, λ) . Thus $\psi(z, \lambda) = 0$ and by (12) if $\nabla_{\mathbb{T}}\psi_c(z, \lambda) = \mathbf{0}$, then $\nabla_{\mathbb{T}}f_p(z, \lambda) = \mathbf{0}$.
- (iv) Exactly one polynomial f_p and one polynomial g_r vanish at (z, λ) . Thus $\psi(z, \lambda) = 0$ and by (12) if $\nabla_{\mathbb{T}}\psi_c(z, \lambda) = \mathbf{0}$, then, after reindexing so that $p = r = 1$, we have

$$(13) \quad \nabla_{\mathbb{T}}f_1(z, \lambda) \cdot \prod_{i=2}^m f_i(z, \lambda) - (-1)^m \nabla_{\mathbb{T}}g_1(z, \lambda) \cdot \prod_{i=2}^m g_i(z, \lambda) = \mathbf{0}.$$

- (v) No polynomials f_i or g_i vanish at (z, λ) .

In each case, we will show that the set of parameters $c \in Y'$ such that there exist (z, λ) satisfying these conditions lies in a proper subvariety of Y' . Cases (i)–(iv) use arguments based on the dimension of fibers and images of a map and are proven in the rest of this section. Case (v) is proven in Section 4.3 and it uses Bertini's Theorem. \square

Let us write X for the space $(\mathbb{C}^\times)^{d+1}$ and x for a point $(z, \lambda) \in X$. We first derive consequences of some vanishing statements. For a finite set $\mathcal{F} \subset \mathbb{Z}^{d+1}$, let $\mathbb{C}^{\mathcal{F}}$ be the space of coefficients of polynomials in $x \in X$ with support \mathcal{F} . This is the parameter space for polynomials with support \mathcal{F} .

Lemma 4.5. *We have the following.*

- (1) For any $x \in X$, $f(x) = 0$ is a nonzero homogeneous linear equation on $\mathbb{C}^{\mathcal{F}}$.
- (2) For any $x \in X$, $\{\nabla_{\mathbb{T}}f(x) \mid f \in \mathbb{C}^{\mathcal{F}}\}$ is the linear span $\mathbb{C}\mathcal{F}$ of \mathcal{F} .

Suppose that the affine span of \mathcal{F} does not contain the origin. Then

- (3) For any $f \in \mathbb{C}^{\mathcal{F}}$ and $x \in X$, $\nabla_{\mathbb{T}}f = \mathbf{0}$ implies that $f(x) = 0$.
- (4) For any $x \in X$, the equation $\nabla_{\mathbb{T}}f(x) = \mathbf{0}$ defines a linear subspace of $\mathbb{C}^{\mathcal{F}}$ of codimension $\dim \mathbb{C}\mathcal{F}$.

Proof. Writing $f = \sum_{a \in \mathcal{F}} c_a x^a$, the first statement is obvious. We have $\nabla_{\mathbb{T}}f = \sum a c_a x^a$. As the coefficients c_a are independent complex numbers and $x^a \neq 0$, Statement (2) is immediate. The hypothesis that the affine span of \mathcal{F} does not contain the origin implies

that any $f \in \mathbb{C}^{\mathcal{F}}$ is quasi-homogeneous. Statement (3) follows from Equation (11). The last statement follows from the observation that the set of f such that $\nabla_{\mathbb{T}} f = \mathbf{0}$ is the kernel of a surjective linear map $\mathbb{C}^{\mathcal{F}} \rightarrow \mathbb{C}^{\mathcal{F}}$. \square

Let $\mathcal{F} := G \cap (\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})$, where $\mathbf{e} = (\mathbf{0}, 1)$, be the (common) support of the diagonal polynomials f_i and let $\mathcal{G} := G \cap \mathcal{A}(\Gamma)$ be the (common) support of the polynomials g_j . We either have that $\mathcal{F} = \mathcal{G}$ or $\mathcal{F} = \mathcal{G} \cup \{\mathbf{e}\}$. Also, $|\mathcal{F}| > 1$ as G is not a vertex, and as G is a proper face of $Q = \text{conv}(\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})$, but not its base, the polynomials f_i, g_j are quasi-homogeneous with a common quasi-homogeneity.

The parameter space for the entries of $(L_c(z) - \lambda I)|_G$ is

$$Z := (\mathbb{C}^{\mathcal{F}})^{\oplus m} \oplus (\mathbb{C}^{\mathcal{G}})^{\oplus m}.$$

We write $c = (f_{\bullet}, g_{\bullet}) = (f_1, \dots, f_m, g_1, \dots, g_m)$ for points of Z . This is a coordinate subspace of the parameter space Y' . As Z contains exactly those parameters that can appear in the facial polynomial $\psi_c(x)$, it suffices to show that the set of parameters $c = (f_{\bullet}, g_{\bullet}) \in Z$ such that $\text{Var}(\psi_c(x))$ is singular lies in a proper subvariety of Z . The same case distinctions (i)–(v) in the proof of Theorem 4.4 apply.

After reindexing, Case (i) in the proof of Theorem 4.4 follows from the next lemma.

Lemma 4.6. *The set*

$$\Theta := \{c \in Z \mid \exists x \in X \text{ with } f_1(x) = f_2(x) = g_1(x) = g_2(x) = 0\}$$

lies in a proper subvariety of Z .

Proof. Consider the incidence correspondence,

$$\Upsilon := \{(x, f_{\bullet}, g_{\bullet}) \in X \times Z \mid f_1(x) = f_2(x) = g_1(x) = g_2(x) = 0\}.$$

This has projections to X and to Z and its image in Z is the set Θ .

Consider the projection $\pi_X: \Upsilon \rightarrow X$. By Lemma 4.5(1), for $x \in X$, each condition $f_i(x) = 0$, $g_i(x) = 0$ for $i = 1, 2$ is a linear equation on $\mathbb{C}^{\mathcal{F}}$ or $\mathbb{C}^{\mathcal{G}}$. These are independent on Z as they involve different variables. Thus the fiber $\pi_X^{-1}(x)$ is a vector subspace of Z of codimension 4, and $\dim(\Upsilon) = \dim(Z) - 4 + \dim(X) = \dim(Z) + d - 3$.

Consider the projection π_Z to Z and let $(f_{\bullet}, g_{\bullet}) \in \pi_Z(\Upsilon)$. Then there is an $x \in X$ such that $f_1(x) = f_2(x) = g_1(x) = g_2(x) = 0$. Let $w \in \mathbb{Z}^{d+1}$ be a common quasi-homogeneity of the polynomials f_i, g_j . By Lemma 3.6 (1), for any $t \in \mathbb{C}^{\times}$, each of f_1, f_2, g_1, g_2 vanishes at $t^w \cdot x$. Thus the fiber $\pi_Z^{-1}(f_{\bullet}, g_{\bullet})$ has dimension at least one. By the Theorem [29, Theorem 1.25] on the dimension of the image and fibers of a map, the image $\pi_Z(\Upsilon)$ has dimension at most $\dim(Z) + d - 4 < \dim(Z)$, which establishes the lemma. \square

After reindexing and possibly interchanging f with g , Cases (ii) and (iii) in the proof of Theorem 4.4 follow from the next lemma.

Lemma 4.7. *The set*

$$\Theta := \{c \in Z \mid \exists x \in X \text{ with } f_1(x) = f_2(x) = g_1(x) = 0 \text{ and } \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}\}$$

lies in a proper subvariety of Z .

Proof. Consider the incidence correspondence,

$$\Upsilon := \{(x, f_\bullet, g_\bullet) \in X \times Z \mid f_1(x) = f_2(x) = g_1(x) = 0 \text{ and } \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}\}.$$

Let $x \in X$ and consider the fiber $\pi_X^{-1}(x)$. As in the proof of Lemma 4.6, the conditions $f_1(x) = f_2(x) = 0$ are two independent linear equations on Z . By Lemma 4.5 (3), $\nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$ implies that $g_1(x) = 0$, and by Lemma 4.5 (4), the condition $\nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$ is $\dim \mathbb{C}\mathcal{G}$ further independent linear equations on Z .

If $|\mathcal{G}| = 1$, so that $g_1 = c_a x^a$ is a single term, then $g(x) = 0$ implies that $c_a = 0$. Consequently, the image Θ of Υ in Z lies in a proper subvariety. Otherwise, $|\mathcal{G}| > 1$ which implies that $\dim \mathbb{C}\mathcal{G} \geq 2$, and thus the fiber has codimension at least 4. As in the proof of Lemma 4.6, this implies that Θ lies in a proper subvariety of Z . \square

Case (iv) in the proof of Theorem 4.4 is more involved.

Lemma 4.8. *The set*

$$\Theta := \{c \in Z \mid \exists x \in X \text{ with } f_1(x) = g_1(x) = 0 \text{ and } \nabla_{\mathbb{T}} \psi_c(x) = \mathbf{0}\}$$

lies in a proper subvariety of Z .

Proof. The set Θ includes the sets of Lemmas 4.6 and 4.7. Let $\Theta^\circ \subset \Theta$ be the set of $c = (f_\bullet, g_\bullet)$ that have a witness $x \in X$ ($f_1(x) = g_1(x) = 0$ and $\nabla_{\mathbb{T}} \psi_c(x) = \mathbf{0}$) such that none of $\nabla_{\mathbb{T}} f_1(x)$, $\nabla_{\mathbb{T}} g_1(x)$, or $f_i(x)g_i(x)$ for $i > 1$ vanish. It will suffice to show that Θ° lies in a proper subvariety of Z .

For this, we use the incidence correspondence,

$$\Upsilon := \{(y, x, f_\bullet, g_\bullet) \in \mathbb{C}^\times \times X \times Z \mid f_1(x) = g_1(x) = 0, \\ y \prod_{i=2}^m f_i(x) - (-1)^m \prod_{i=2}^m g_i(x) = 0, \text{ and } \nabla_{\mathbb{T}} f_1(x) - (-1)^m y \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}\}.$$

We show that $\Theta^\circ \subset \pi_Z(\Upsilon)$. Let $c = (f_\bullet, g_\bullet) \in \Theta^\circ$ with witness $x \in X$ in that $f_1(x) = g_1(x) = 0$ and $\nabla_{\mathbb{T}} \psi_c(x) = \mathbf{0}$, but none of $\nabla_{\mathbb{T}} f_1(x)$, $\nabla_{\mathbb{T}} g_1(x)$, or $f_i(x)g_i(x)$ for $i > 1$ vanish. There is a unique $y \in \mathbb{C}^\times$ satisfying

$$y \prod_{i=2}^m f_i(x) - (-1)^m \prod_{i=2}^m g_i(x) = 0.$$

Dividing (13) by $\prod_{i=2}^m f_i(x)$ gives $\nabla_{\mathbb{T}} f_1(x) - (-1)^m y \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$, and thus $(y, x, f_\bullet, g_\bullet) \in \Upsilon$.

We now determine the dimension of Υ . Let $(y, x) \in \mathbb{C}^\times \times X$ and consider the fiber $\pi^{-1}(y, x) \subset Z$ above it in Υ . The two linear and one nonlinear equations

$$(14) \quad f_1(x) = g_1(x) = y \prod_{i=2}^m f_i(x) - (-1)^m \prod_{i=2}^m g_i(x) = 0$$

are independent on Z as they involve disjoint sets of variables, and thus define a subvariety $T \subset Z$ of codimension 3. Consider the remaining equation, $\nabla_{\mathbb{T}} f_1(x) - (-1)^m y \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$.

Note that if $\mathbf{e} = (\mathbf{0}, 1)$ lies in the support \mathcal{F} of f_1 , so that $\mathcal{F} = \mathcal{G} \cup \{\mathbf{e}\}$, then $\nabla_{\mathbb{T}} f_1(x)$ contains the term $-\mathbf{e}$ and thus cannot lie in the span $\mathbb{C}\mathcal{G}$ of \mathcal{G} , which contains $\nabla_{\mathbb{T}} g_1(x)$ by Lemma 4.5(2). In this case the fiber is empty and $\Theta^\circ = \emptyset$.

Suppose that $\mathcal{F} = \mathcal{G}$ and $(f_\bullet, g_\bullet) \in T$. Let $w \in \mathbb{Z}^{d+1}$ be any homogeneity for f_1 (or g_1). Then there exists $w_{\mathcal{F}} \neq 0$ such that $w \cdot a = w_{\mathcal{F}}$ for all $a \in \mathcal{F}$. Equation (11) implies that

$$w \cdot \nabla_{\mathbb{T}} f_1(x) = w_{\mathcal{F}} f_1(x) = 0,$$

and the same for g_1 . Thus $\nabla_{\mathbb{T}} f_1(x)$ and $\nabla_{\mathbb{T}} g_1(x)$ are annihilated by all homogeneities and so lie in the affine span of \mathcal{F} —the linear span of differences $a-b$ for $a, b \in \mathcal{F}$. This has dimension $\dim \mathbb{C}\mathcal{F} - 1$. Consequently, $\nabla_{\mathbb{T}} f_1(x) - (-1)^m y \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$ consists of $\dim \mathbb{C}\mathcal{F} - 1$ independent linear equations on the subset of $\mathbb{C}^{\mathcal{F}} \oplus \mathbb{C}^{\mathcal{F}}$ consisting of pairs f_1, g_1 such that $f_1(x) = g_1(x) = 0$. These are independent of the third equation in (14). Thus the fiber $\pi^{-1}(y, x) \subset Z$ has codimension $3 + \dim \mathbb{C}\mathcal{F} - 1 = 2 + \dim \mathbb{C}\mathcal{F}$ and so

$$\dim \Upsilon = \dim(\mathbb{C}^\times \times X) + \dim Z - \dim \mathbb{C}\mathcal{F} - 2 = \dim Z + d - \dim \mathbb{C}\mathcal{F}.$$

Let $(f_\bullet, g_\bullet) \in \pi_Z(\Upsilon)$ have witness (y, x) . That is, the equations (14) hold, as well as $\nabla_{\mathbb{T}} f_1(x) - (-1)^m y \nabla_{\mathbb{T}} g_1(x) = \mathbf{0}$. As in the proof of Lemma 4.6, if $w \in \mathbb{Z}^{d+1}$ is a quasi-homogeneity for polynomials of support \mathcal{F} , then $(y, t^w \cdot x)$ also satisfies these equations.

We have $\mathcal{F} = \mathcal{G} = G \cap \mathcal{A}(\Gamma)$, so that G is a face of the base of Q . Thus there are at least two (in fact the codimension of G in Q) independent homogeneities, which implies that the fiber $\pi_Z^{-1}(f_\bullet, g_\bullet)$ has dimension at least two. This implies that the image Θ° has dimension at most $\dim Z + d - \dim \mathbb{C}\mathcal{F} - 2$. Since G is not a vertex, $\dim \mathbb{C}\mathcal{F} \geq 2$, which shows that $\dim \Theta^\circ < \dim Z$ and completes the proof. \square

4.3. Case (v). For $\alpha \in \mathbb{C}^\times$, define $\Psi(\alpha, f_\bullet, g_\bullet) \subset X$ to be the set

$$\left\{ x \in X \mid \text{none of } f_i(x)g_i(x) \text{ for } i \geq 1 \text{ vanish and } \prod_{i=1}^m f_i(x) - (-1)^m \alpha \prod_{i=1}^m g_i(x) = 0 \right\}.$$

Case (v) in the proof of Theorem 4.4 follows from the next lemma.

Lemma 4.9. *There is a dense open subset $U_1 \subset Z$ such that if $(f_\bullet, g_\bullet) \in U_1$, then $\Psi(1, f_\bullet, g_\bullet)$ is smooth.*

We will deduce this from a weaker lemma.

Lemma 4.10. *There is a dense open subset $U \subset \mathbb{C}^\times \times Z$ such that if $(\alpha, f_\bullet, g_\bullet) \in U$, then $\Psi(\alpha, f_\bullet, g_\bullet)$ is smooth.*

Proof of Lemma 4.9. If we knew that the set U of Lemma 4.10 contained a point $(1, f_\bullet, g_\bullet)$, then $U_1 := U \cap (\{1\} \times Z)$ would be a dense open subset of Z , which would complete the proof. As we do not know this, we must instead argue indirectly.

Suppose that there is no such open set U_1 as in Lemma 4.9. Then the set $\Xi \subset Z$ consisting of (f_\bullet, g_\bullet) such that $\Psi(1, f_\bullet, g_\bullet)$ is singular is dense in Z .

For $\alpha \in \mathbb{C}^\times$ and $(f_\bullet, g_\bullet) \in Z$, define $\alpha.(f_\bullet, g_\bullet)$ to be $(f_\bullet, \alpha.g_\bullet)$ where

$$\alpha.(g_1, g_2, \dots, g_m) = (\alpha g_1, g_2, \dots, g_m).$$

This is a \mathbb{C}^\times -action on Z . Consequently, $\alpha.\Xi$ is dense in Z for all $\alpha \in \mathbb{C}^\times$.

Let $U \subset \mathbb{C}^\times \times Z$ be the set of Lemma 4.10. As it is nonempty, let $(\alpha, f'_\bullet, g'_\bullet) \in U$. Then $U_\alpha := U \cap (\{\alpha\} \times Z)$ is nonempty and open in $\{\alpha\} \times Z$. As $\alpha.\Xi$ is dense, we have

$$U_\alpha \cap (\{\alpha\} \times \alpha.\Xi) \neq \emptyset.$$

This is a contradiction, for if $(\alpha, f_\bullet, g_\bullet) \in U_\alpha$, then $\Psi(\alpha, f_\bullet, g_\bullet)$ is smooth, but if $(f_\bullet, g_\bullet) \in \alpha.\Xi$, then $(f_\bullet, \alpha^{-1}g_\bullet) \in \Xi$ and $\Psi(1, f_\bullet, \alpha^{-1}g_\bullet)$ is singular. The contradiction follows from the equality of sets $\Psi(\alpha, f_\bullet, g_\bullet) = \Psi(1, f_\bullet, \alpha^{-1}g_\bullet)$. \square

Proof of Lemma 4.10. Let $T \subset X \times Z$ be the set of $(x, f_\bullet, g_\bullet)$ such that none of $f_i(x)g_i(x)$ for $i \geq 1$ vanish. Define $\varphi: T \rightarrow \mathbb{C}^\times \times Z$ by

$$\varphi(x, f_\bullet, g_\bullet) = \left((-1)^m \prod_{i=1}^m f_i(x) / \prod_{i=1}^m g_i(x), f_\bullet, g_\bullet \right).$$

Notice that $\varphi^{-1}(\alpha, f_\bullet, g_\bullet) = \Psi(\alpha, f_\bullet, g_\bullet)$ for $(\alpha, f_\bullet, g_\bullet) \in \mathbb{C}^\times \times Z$.

We claim that $\varphi(T)$ is dense in $\mathbb{C}^\times \times Z$. For this, recall that the polynomials f_i have support \mathcal{F} , which is $G \cap (\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})$ for some face G of $Q = \text{conv}(\mathcal{A}(\Gamma) \cup \{\mathbf{e}\})$ that is neither its base nor a vertex, and the polynomials g_i have support $\mathcal{G} = G \cap \mathcal{A}(\Gamma)$. Since G is not a vertex, there are $a, b \in \mathcal{F}$ with $a \neq b$ and $b \in \mathcal{A}(\Gamma)$.

Let $f_i := x^a$ and $g_i := x^b$ for $i = 1, \dots, m$. Then $X \times \{(f_\bullet, g_\bullet)\} \subset T$ and for $x \in X$ $\varphi(x, f_\bullet, g_\bullet) = (x^{ma} - (-1)^m x^{mb}, f_\bullet, g_\bullet)$. The map $X = (\mathbb{C}^\times)^{d+1} \rightarrow \mathbb{C}^\times$ given by $x \mapsto x^{ma} - (-1)^m x^{mb}$ is surjective as $ma - mb \neq 0$. This implies that the differential $d\varphi$ is surjective at any point of $X \times \{(f_\bullet, g_\bullet)\}$, and therefore $\varphi(T)$ is dense in $\mathbb{C}^\times \times Z$.

Since T is an open subset of the smooth variety $X \times Z$, it is smooth. Then Bertini's Theorem [29, Thm. 2.27, p. 139] implies that there is a dense open subset $U \subset \mathbb{C}^\times \times Z$ such that for $(\alpha, f_\bullet, g_\bullet) \in U$, $\varphi^{-1}(\alpha, f_\bullet, g_\bullet) = \Psi(\alpha, f_\bullet, g_\bullet)$ is smooth. \square

5. CRITICAL POINTS PROPERTY

We illustrate our results, using them to establish the critical points property (and thus the spectral edges nondegeneracy conjecture) for three periodic graphs. We first state this property.

Let Γ be a connected \mathbb{Z}^d -periodic graph with parameter space $Y = \mathbb{C}^E \times \mathbb{C}^W$ for discrete operators on Γ . We say that Γ has the *critical points property* if there is a dense open subset $U \subset Y$ such that if $c \in U$, then every critical point of the function λ on the Bloch variety $\text{Var}(D_c(z, \lambda))$ is nondegenerate in that the Hessian determinant

$$(15) \quad \det \left(\left(\frac{\partial^2 \lambda}{\partial z_i \partial z_j} \right)_{i,j=1}^d \right)$$

is nonzero at that critical point. Here, the derivatives are implicit, using that $D(z, \lambda) = 0$.

5.1. Reformulation of Hessian condition. Let $D = \det(L_c(z) - \lambda I)$ be the dispersion function for an operator L_c on a periodic graph Γ . In Section 2 we derived the equations for the critical points of the function λ on the Bloch variety $\text{Var}(D(z, \lambda))$,

$$(16) \quad D(z, \lambda) = 0 \quad \text{and} \quad \frac{\partial D}{\partial z_i} = 0 \quad \text{for } i = 1, \dots, d.$$

Implicit differentiation of $D = 0$ gives $\frac{\partial D}{\partial z_j} + \frac{\partial D}{\partial \lambda} \cdot \frac{\partial \lambda}{\partial z_j} = 0$. If $\frac{\partial D}{\partial \lambda} \neq 0$, then $\frac{\partial \lambda}{\partial z_j} = 0$. If $\frac{\partial D}{\partial \lambda} = 0$, then (z, λ) is a singular point hence is also a critical point of the function λ and so we again

have $\frac{\partial \lambda}{\partial z_j} = 0$. Differentiating again we obtain,

$$0 = \frac{\partial}{\partial z_i} \left(\frac{\partial D}{\partial z_j} + \frac{\partial D}{\partial \lambda} \cdot \frac{\partial \lambda}{\partial z_j} \right) = \frac{\partial^2 D}{\partial z_i \partial z_j} + \frac{\partial^2 D}{\partial z_i \partial \lambda} \cdot \frac{\partial \lambda}{\partial z_j} + \frac{\partial D}{\partial \lambda} \cdot \frac{\partial^2 \lambda}{\partial z_i \partial z_j}.$$

At a critical point (so that $\frac{\partial \lambda}{\partial z_j} = 0$), we have

$$\frac{\partial^2 D}{\partial z_i \partial z_j} = -\frac{\partial D}{\partial \lambda} \cdot \frac{\partial^2 \lambda}{\partial z_i \partial z_j}.$$

Thus

$$\det \left(\left(\frac{\partial^2 D}{\partial z_i \partial z_j} \right)_{i,j=1}^d \right) = \left(-\frac{\partial D}{\partial \lambda} \right)^d \cdot \det \left(\left(\frac{\partial^2 \lambda}{\partial z_i \partial z_j} \right)_{i,j=1}^d \right).$$

Consider now the Jacobian matrix of the critical point equations (16),

$$J = \begin{pmatrix} \frac{\partial D}{\partial z_1} & \cdots & \frac{\partial D}{\partial z_d} & \frac{\partial D}{\partial \lambda} \\ \frac{\partial^2 D}{\partial z_1^2} & \cdots & \frac{\partial^2 D}{\partial z_d \partial z_1} & \frac{\partial^2 D}{\partial \lambda \partial z_1} \\ \vdots & \ddots & \vdots & \vdots \\ \frac{\partial^2 D}{\partial z_1 \partial z_d} & \cdots & \frac{\partial^2 D}{\partial z_d^2} & \frac{\partial^2 D}{\partial \lambda \partial z_d} \end{pmatrix}.$$

At a critical point, the first row is $(0 \cdots 0 \frac{\partial D}{\partial \lambda})$, and thus

$$\det(J) = \frac{\partial D}{\partial \lambda} \det \left(\left(\frac{\partial^2 D}{\partial z_i \partial z_j} \right)_{i,j=1}^d \right) = (-1)^d \left(\frac{\partial D}{\partial \lambda} \right)^{d+1} \cdot \det \left(\left(\frac{\partial^2 \lambda}{\partial z_i \partial z_j} \right)_{i,j=1}^d \right).$$

A solution ζ of a system of polynomial equations on \mathbb{C}^n is *regular* if the Jacobian of the system at ζ has full rank n . Regular solutions are isolated and have multiplicity 1. We deduce the following lemma.

Lemma 5.1. *A nonsingular critical point (z, λ) on $\text{Var}(D_c(z, \lambda))$ is nondegenerate if and only if it is a regular solution of the critical point equations (16).*

The following theorem is adapted from arguments in [9, Sect. 5.4].

Theorem 5.2. *Let Γ be a \mathbb{Z}^d -periodic graph. If there is a parameter value $c \in Y$ such that the critical point equations have $(d+1)!\text{vol}(\mathcal{N}(\Gamma))$ regular solutions, then the critical points property holds for Γ .*

Proof. Let Y be the parameter space for operators L on Γ . Consider the variety

$$CP := \{(c, z, \lambda) \in Y \times (\mathbb{C}^\times)^d \times \mathbb{C} \mid \text{the critical point equations (8) hold}\},$$

which is the incidence variety of critical points on all Bloch varieties for operators on Γ . Let π be its projection to Y . For any $c \in Y$, the fiber $\pi^{-1}(c)$ is the set of critical points of the function λ on the corresponding Bloch variety for D_c . By Corollary 2.5, there are at most $(d+1)!\text{vol}(\mathcal{N}(D_c))$ isolated points in the fiber.

Let $c \in Y$ be a point such that the critical point equations have $(d+1)!\text{vol}(\mathcal{N}(\Gamma))$ regular solutions. Then $(d+1)!\text{vol}(\mathcal{N}(\Gamma)) \leq (d+1)!\text{vol}(\mathcal{N}(D_c))$. By Lemma 4.3, $\mathcal{N}(D_c)$ is a subpolytope of $\mathcal{N}(\Gamma)$, so that $\text{vol}(\mathcal{N}(D_c)) \leq \text{vol}(\mathcal{N}(\Gamma))$. We conclude that both polytopes have

the same volume and are therefore equal. In particular, the corresponding Bloch variety has the maximum number of critical points, and each is a regular solution of the critical point equations (8). Because they are regular solutions, the implicit function theorem implies that there is a neighborhood U_c of c in the classical topology on Y such that the map $\pi^{-1}(U_c) \rightarrow U_c$ is proper (it is a $(d+1)!\text{vol}(\mathcal{N}(\Gamma))$ -sheeted cover).

The set DC of degenerate critical points is the closed subset of CP given by the vanishing of the Hessian determinant (15). Since π is proper over U_c , if $DP = \pi(DC)$ is the image of DC in Y , then $DP \cap U_c$ is closed in U_c . As the points of $\pi^{-1}(c)$ are regular solutions, Lemma 5.1 implies they are all nondegenerate and thus $c \notin DP$, so that $U_c \setminus DP$ is a nonempty classically open subset of Y consisting of parameter values c' with the property that all critical points on the corresponding Bloch variety are nondegenerate.

This implies that there is a nonempty Zariski open subset of Y consisting of parameters such that all critical points on the corresponding Bloch variety are nondegenerate, which completes the proof. \square

By Theorem 5.2, it suffices to find a single Bloch variety with the maximum number of isolated critical points to establish the critical points property for a periodic graph. The following examples use such a computation to establish the critical points property for $2^{19}+2$ graphs Γ . Computer code and output are available at the github repository².

Example 5.3. Let us consider the dense \mathbb{Z}^2 -periodic graph Γ of Figure 6. It has $m = 2$ points in its fundamental domain and the convex hull of its support $\mathcal{A}(\Gamma)$ has area 4. By Theorem C, a general operator on Γ has $2! \cdot 2^{2+1} \cdot 4 = 64$ critical points. There are 13 edges




FIGURE 8. Dense periodic graph and its polytope from Figure 6.

and two vertices in W , and independent computations in the computer algebra systems Macaulay2 [17] and Singular [8] find a point $c \in Y = \mathbb{C}^{15}$ such that the critical point equations have 64 regular solutions on $(\mathbb{C}^\times)^2 \times \mathbb{C}$. By Theorem 5.2, the critical points property holds for Γ . These computations are independent in that the code, authors, and parameter values for each are distinct. \diamond

²<https://mattfaust.github.io/CPDPO>.

Example 5.4. The graph Γ in Figure 9 is not dense. Its restriction to the fundamental domain is not the complete graph on 3 vertices and there are three and not nine edges between any two adjacent translates of the fundamental domain. Altogether, it has $3 \cdot 6 + 1 = 19$ fewer edges than the corresponding dense graph. Its support $\mathcal{A}(\Gamma)$ forms the columns of the matrix $\begin{pmatrix} 0 & 1 & 1 & 0 & -1 & -1 & 0 \\ 0 & 0 & 1 & 1 & 0 & -1 & -1 \end{pmatrix}$ whose convex hull is a hexagon of area 3.




FIGURE 9. Sparse graph with the same Newton polytope as the corresponding dense graph.

Despite Γ not being dense, its Newton polytope $\mathcal{N}(\Gamma)$ is equal to the Newton polytope of the dense graph with the same parameters, $\mathcal{A}(\Gamma)$ and W . Figure 9 displays the Newton polytope, along with elements of the support of the dispersion function that are visible. Observe that on each triangular face, there are four and not ten monomials.

By Theorem A (Corollary 2.5), there are at most $2! \cdot 3^{2+1} \cdot 3 = 162$ critical points. There are eleven edges and three vertices in W , and independent computations in Macaulay2 and Singular find a point $c \in Y = \mathbb{C}^{14}$ such that the critical point equations have 162 regular solutions on $(\mathbb{C}^\times)^2 \times \mathbb{C}$. By Theorem 5.2, the critical points property holds for Γ .

Let Γ' be a graph that has the same vertex set and support as Γ , and contains all the edges of Γ —then [9, Thm. 22] implies that the critical points property also holds for Γ' . This establishes the critical points property for an additional $2^{19} - 1$ periodic graphs. \diamond

Example 5.5. The graph Γ of Figure 10 has only ten edges but the same fundamental domain W and support $\mathcal{A}(\Gamma)$ as the the graph of Figure 9, which had eleven edges. Its Newton polytope is smaller, as it is missing the vertices $(3, 3, 0)$ and $(-3, -3, 0)$.

It has volume $70/3$ and normalized volume $3! \cdot 70/3 = 140$. Independent computations in Macaulay2 and Singular find a point $c \in Y = \mathbb{C}^{13}$ such that the critical point equations have 140 regular solutions on $(\mathbb{C}^\times)^2 \times \mathbb{C}$. Thus there are no critical points at infinity, and Theorem B implies that the Bloch variety is smooth at infinity.

As before, achieving the bound of Corollary 2.5 with regular solutions implies that all critical points are nondegenerate and the critical points property holds for Γ . \diamond




FIGURE 10. A periodic graph and its Newton polytope.

6. CONCLUSION

We considered the critical points of the complex Bloch variety for an operator on a periodic graph. We gave a bound on the number of critical points—the normalized volume of a Newton polytope—together with a criterion for when that bound is attained. We presented a class of graphs (dense periodic graphs) and showed that this criterion holds for general discrete operators on a dense graph. Lastly, we used these results to find $2^{19} + 2$ graphs on which the spectral edges conjecture holds for general discrete operators when $d = 2$.

REFERENCES

1. G. Berkolaiko, Y. Canzani, G. Cox, and J.L. Marzuola, A local test for global extrema in the dispersion relation of a periodic graph, *Pure and Applied Analysis* **4** (2022), no. 2, 257–286.
2. G. Berkolaiko and P. Kuchment, Introduction to quantum graphs, *Mathematical Surveys and Monographs*, vol. 186, American Mathematical Society, Providence, RI, 2013.
3. D. N. Bernstein, The number of roots of a system of equations, *Functional Anal. Appl.* **9** (1975), no. 3, 183–185.
4. P. Breiding, F. Sottile, and J. Woodcock, Euclidean distance degree and mixed volume, *Found. Comput. Math.* (2021).
5. Y. Colin de Verdière, Sur les singularités de van Hove génériques, *Analyse globale et physique mathématique* (Lyon, 1989), no. 46, Soc. Math. France, 1991, pp. 99–110.
6. D. Cox, J. Little, and D. O’Shea, Using algebraic geometry, second ed., *Graduate Texts in Mathematics*, vol. 185, Springer, New York, 2005.
7. ———, Ideals, varieties, and algorithms, third ed., *Undergraduate Texts in Mathematics*, Springer, New York, 2007.
8. W. Decker, G-M. Greuel, G. Pfister, and H. Schönemann, SINGULAR 4-3-0 — A computer algebra system for polynomial computations, <http://www.singular.uni-kl.de>, 2022.
9. N. Do, P. Kuchment, and F. Sottile, Generic properties of dispersion relations for discrete periodic operators, *J. Math. Phys.* **61** (2020), no. 10, 103502, 19.
10. G. Ewald, Combinatorial convexity and algebraic geometry, *Graduate Texts in Mathematics*, vol. 168, Springer-Verlag, New York, 1996.
11. J. Fillman, W. Liu, and R. Matos, Irreducibility of the Bloch variety for finite-range Schrödinger operators, *Journal of Functional Analysis* **283** (2022), no. 10, 109670.

12. ———, Algebraic properties of the Fermi variety for periodic graph operators, 2023, [arXiv:2305.06471](https://arxiv.org/abs/2305.06471).
13. N. Filonov and I. Kachkovskiy, On the structure of band edges of 2-dimensional periodic elliptic operators, *Acta Math.* **221** (2018), no. 1, 59–80. MR 3877018
14. Wm. Fulton, Intersection theory, second ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge.*, vol. 2, Springer-Verlag, Berlin, 1998.
15. I. M. Gel'fand, M. M. Kapranov, and A. V. Zelevinsky, Discriminants, resultants, and multidimensional determinants, *Mathematics: Theory & Applications*, Birkhäuser, Boston, MA, 1994.
16. D. Gieseker, H. Knörrer, and E. Trubowitz, The geometry of algebraic Fermi curves, *Perspectives in Mathematics*, vol. 14, Academic Press, Inc., Boston, MA, 1993.
17. D.R. Grayson and M.E. Stillman, Macaulay2, a software system for research in algebraic geometry, Available at <http://www.math.uiuc.edu/Macaulay2/>.
18. F. Klopp and J. Ralston, Endpoints of the spectrum of periodic operators are generically simple, *Methods Appl. Anal.* **7** (2000), no. 3, 459–463.
19. A. G. Kouchnirenko, Polyèdres de Newton et nombres de Milnor, *Invent. Math.* **32** (1976), no. 1, 1–31.
20. C. Kravaris, On the density of eigenvalues on periodic graphs, *SIAM Journal on Applied Algebra and Geometry*, to appear. [arXiv.org/2103:12734](https://arxiv.org/abs/2103.12734), 2021.
21. P. Kuchment, Floquet theory for partial differential equations, *Operator Theory: Advances and Applications*, vol. 60, Birkhäuser Verlag, Basel, 1993.
22. ———, An overview of periodic elliptic operators, *Bull. Amer. Math. Soc. (N.S.)* **53** (2016), no. 3, 343–414.
23. W. Li and S.P. Shipman, Irreducibility of the Fermi surface for planar periodic graph operators, *Lett. Math. Phys.* **110** (2020), no. 9, 2543–2572.
24. J. Lindberg, N. Nicholson, J.L. Rodriguez, and Z. Wang, The maximum likelihood degree of sparse polynomial systems, *SIAM Journal on Applied Algebra and Geometry* **7** (2023), no. 1.
25. W. Liu, Irreducibility of the Fermi variety for discrete periodic Schrödinger operators and embedded eigenvalues, *Geom. Funct. Anal.* **32** (2022), no. 1, 1–30.
26. ———, Fermi isospectrality of discrete periodic Schrödinger operators with separable potentials on \mathbb{Z}^2 , 2023, pp. 1139–1149. MR 4576766
27. S. P. Novikov, Bloch functions in the magnetic field and vector bundles. Typical dispersion relations and their quantum numbers, *Dokl. Akad. Nauk SSSR* **257** (1981), no. 3, 538–543.
28. ———, Two-dimensional Schrödinger operators in periodic fields, *Current problems in mathematics*, Vol. 23, *Itogi Nauki i Tekhniki*, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1983, pp. 3–32.
29. I. R. Shafarevich, Basic algebraic geometry 1, translated from the 2007 third russian edition ed., Springer, Heidelberg, 2013.
30. F. Sottile, Real solutions to equations from geometry, *University Lecture Series*, vol. 57, American Mathematical Society, Providence, RI, 2011.
31. B. Sturmfels, Gröbner bases and convex polytopes, *University Lecture Series*, vol. 8, American Mathematical Society, Providence, RI, 1996.

MATTHEW FAUST, DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843, USA

E-mail address: mfaust@tamu.edu

URL: <http://mattfaust.github.io>

FRANK SOTTILE, DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843, USA

E-mail address: sottile@tamu.edu

URL: <https://franksottile.github.io>