

Frontiers of Reality in Schubert Calculus

Mathematics Advisory Board of the Lorentz Center


10 June 2009, Lorentz Center, Leiden



Frank Sottile

sottile@math.tamu.edu

In which the theory of integrable systems resolves a compelling conjecture in the real Schubert calculus by reducing it to an elementary fact from linear algebra.



Polynomial systems with only real roots

Among the roots of a real univariate polynomial f , some are real and the rest occur in complex-conjugate pairs.

Rarely are all roots of f real.

A primary example that comes to mind is when f is the characteristic polynomial of a real symmetric matrix, which only has real eigenvalues.

Similarly, a first example of a system of multivariate polynomials with only real solutions is the system for the eigenvalues/eigenvectors of a symmetric matrix.

It will turn out that this is the elementary fact from linear algebra behind this reality in Schubert calculus.

Wronski map and MTV Theorem

The Wronskian of degree- d polynomials $f_0, \dots, f_n \in \mathbb{C}[t]$ is

$$Wr := \det \begin{pmatrix} f_0(t) & f_1(t) & \cdots & f_n(t) \\ f_0'(t) & f_1'(t) & \cdots & f_n'(t) \\ \vdots & \vdots & \ddots & \vdots \\ f_0^{(n)}(t) & f_1^{(n)}(t) & \cdots & f_n^{(n)}(t) \end{pmatrix}.$$


Up to scalar, Wr depends only on the linear span P of the f_i , and only finitely many spans P have a given Wronskian.

Theorem. (Mukhin, Tarasov, Varchenko) *If $Wr(P)$ has only real roots, then P has a basis of real polynomials.*

More generally, (!) “Geometric problems in Schubert calculus on a Grassmannian involving osculating flags have only real solutions.”

MTV Theorem in 3-space

Let $\gamma(t) = (t, t^2, t^3)$ be the rational normal curve



A cubic polynomial $f(t)$ \iff affine function applied to γ
 \iff affine hyperplane, f^\perp

Two polynomials f, g \iff two affine hyperplanes
 \iff a line $f^\perp \cap g^\perp$

$Wr(f, g)(s) = 0$ \iff the line $f^\perp \cap g^\perp$ meets
tangent line to γ at $\gamma(s)$.

$Wr(f, g)$ is a given
quartic F \iff $f^\perp \cap g^\perp$ meets tangents to γ
at the four roots of F

Numerical accident ?

The proof begins with a numerical accident.

In 1884, Schubert (essentially) determined that

$$\# W r^{-1}(F(t)) = [(n+1)(d-n)]! \frac{1!2! \cdots n!}{(d-n)!(d-n+1)! \cdots d!}.$$

Call this number $\deg(n, d)$.

$\deg(n, d)$ is also the dimension of the space of invariants

$$\left(\left(\mathbb{C}^{n+1} \right)^{\otimes (n+1)(d-n)} \right)^{\mathfrak{sl}_{n+1} \mathbb{C}}.$$

Strengthening this coincidence is at the heart of our story.

A remarkable function

For $i = 0, \dots, n$ let $y_i(t)$ be a polynomial of degree $(i+1)(d-n)$. Define the master function

$$\Phi := \prod_{i=0}^n \text{Discr}(y_i) \Big/ \prod_{i=1}^n \text{Res}(y_{i-1}, y_i) ,$$

where **Discr** and **Res** are the classical discriminant and resultant.

Writing Φ in terms of the roots $s_{i,j}$ of y_i gives,

$$\Phi(s) = \prod_{i=0}^n \prod_{j \neq k} (s_{i,j} - s_{i,k})^2 \cdot \prod_{i=1}^n \prod_{j,k} (s_{i-1,j} - s_{i,k})^{-1} .$$


(The exponents come from the Cartan matrix of type A .)

Remarkably, $\text{deg}(n, d)$ counts the (orbits of) critical points of $\Phi(s)$.


Schematic of proof

Critical points of Φ


Schematic of proof




Schematic of proof




Schematic of proof



Schematic of proof



Schematic of proof



Critical points of master function

For $i = 0, \dots, n$ let $y_i(t)$ be a polynomial of degree $(i+1)(d-n)$. Recall the master function

$$\Phi := \prod_{i=0}^n \text{Discr}(y_i) \Big/ \prod_{i=1}^n \text{Res}(y_{i-1}, y_i) .$$

Fix $y_n(t)$ to be a polynomial of degree $(n+1)(d-n)$ with roots $\mathbf{s} = (s_1, \dots, s_{(n+1)(d-n)})$. This will be our Wronski polynomial.

The master function $\Phi_{\mathbf{s}}(\mathbf{x})$ depends on the roots \mathbf{x} of the other y_i .

Let \mathbf{x} be a critical point of $\Phi_{\mathbf{s}}(\mathbf{x})$, and $\mathbf{y} := (y_0, \dots, y_{n-1})$ the corresponding polynomials whose roots are \mathbf{x} .

Theorem. (MV) *There are $\deg(n, d)$ such critical points \mathbf{y} .*

Spaces of polynomials from \mathbf{y}

For polynomials $\mathbf{y} = (y_0, \dots, y_{n-1})$, with $\deg y_i = (i+1)(d-n)$, the *fundamental differential operator* $D_{\mathbf{y}}$ is

$$\left(\frac{d}{dt} - \ln' \left(\frac{y_n}{y_{n-1}}\right)\right) \cdots \left(\frac{d}{dt} - \ln' \left(\frac{y_1}{y_0}\right)\right) \left(\frac{d}{dt} - \ln'(y_0)\right) .$$

Let $V_{\mathbf{y}}$ be the kernel of $D_{\mathbf{y}}$.

Theorem. (MV)

1. $V_{\mathbf{y}}$ is a space of polynomials iff \mathbf{y} is a critical point of $\Phi_{\mathbf{s}}$.
2. If f_0, \dots, f_n span $V_{\mathbf{y}}$ with $\deg f_i = d-n+i$, then

$$\begin{aligned} y_0 &= f_0, \\ y_1 &= Wr(f_0, f_1), \\ &\vdots \\ y_n &= Wr(f_0, f_1, \dots, f_n). \end{aligned}$$

Periodic Gaudin model

V := dual of vector representation of $\mathfrak{sl}_{n+1}\mathbb{C}$ (also $\mathfrak{gl}_{n+1}\mathbb{C}$). Let $e_{i,j} \in \mathfrak{gl}_{n+1}\mathbb{C}$ be the elementary matrix with 1 in i, j position.

Define an operator $X_{i,j}(t) := \delta_{i,j} \frac{d}{dt} - \sum_{k=1}^{(n+1)(d-n)} \frac{e_{i,j}^{(k)}}{t - s_k}$,

where $e_{i,j}^{(k)}$ acts on the k th factor in $V^{\otimes(n+1)(d-n)}$.

Formal conjugate of the expansion of the **row** determinant of $(X_{i,j})$ is

$$\frac{d^{n+1}}{dt^{n+1}} + K_1(t) \frac{d^n}{dt^n} + \cdots + K_n(t) \frac{d}{dt} + K_{n+1}(t).$$

$K_1(t), \dots, K_{n+1}(t)$ are the **Gaudin Hamiltonians**. They form a family of commuting operators on $V^{\otimes(n+1)(d-n)}$, centralizing $\mathfrak{gl}_{n+1}\mathbb{C}$.

Bethe Ansatz for Gaudin model

In the theory of integrable systems, **Bethe Ansätze** are conjectural methods to find the joint eigenvectors and spectra of families of commuting operators.

As the Gaudin Hamiltonians centralize the action of $\mathfrak{sl}_{n+1}\mathbb{C}$, the Bethe Ansatz also gives a precise way to understand $\left(V^{\otimes(n+1)(d-n)}\right)^{\mathfrak{sl}_{n+1}\mathbb{C}}$.

The idea is to define a (rational) **universal weight function**

$$\beta : \underbrace{y_0, \dots, y_{n-1}}_{\mathbf{x}}, \underbrace{y_n}_{\mathbf{s}} \longrightarrow \text{0-weight space of } V^{\otimes(n+1)(d-n)},$$

and then address for which values (\mathbf{x}, \mathbf{s}) is $\beta(\mathbf{x}, \mathbf{s})$ $\mathfrak{sl}_{n+1}\mathbb{C}$ -invariant.

Completeness of the Bethe Ansatz

Theorem. (MTV) Let \mathbf{x} be a critical point of the master function $\Phi_{\mathbf{s}}$.

1. $\beta(\mathbf{x}, \mathbf{s})$ is well-defined, non-zero, and a joint eigenvector of the Gaudin Hamiltonians.
2. $\beta(\mathbf{x}, \mathbf{s})$ is a $\mathfrak{sl}_{n+1}\mathbb{C}$ -invariant.
3. When \mathbf{s} is general, the vectors $\beta(\mathbf{x}, \mathbf{s})$ for \mathbf{x} a critical point form a basis of $(V^{\otimes(n+1)(d-n)})^{\mathfrak{sl}_{n+1}\mathbb{C}}$.
4. When \mathbf{s} is general, the Gaudin Hamiltonians have simple spectrum.
5. The eigenvalues $\lambda_i(t)$ of $K_i(t)$ on $\beta(\mathbf{x}, \mathbf{s})$ satisfy

$$\frac{d^{n+1}}{dt^{n+1}} + \lambda_1(t) \frac{d^n}{dt^n} + \cdots + \lambda_n(t) \frac{d}{dt} + \lambda_{n+1}(t) = \left(\frac{d}{dt} - \ln' \left(\frac{y_n}{y_{n-1}} \right) \right) \cdots \left(\frac{d}{dt} - \ln' \left(\frac{y_1}{y_0} \right) \right) \left(\frac{d}{dt} - \ln' (y_0) \right),$$

the fundamental differential operator of the critical point \mathbf{x} .

Proof of the Shapiro Conjecture

Usual Euclidean inner product on V induces the [Shapovalov form](#) on $V^{\otimes(n+1)(d-n)}$, which is $\mathfrak{sl}_{n+1}\mathbb{C}$ -invariant.

Gaudin Hamiltonians are symmetric w.r.t the Shapovalov form.

Therefore, when s and t are real, their eigenvalues $\lambda_i(t)$ on a vector $\beta(\mathbf{x}, s)$ for a critical point \mathbf{x} are real.

Then the fundamental differential operator is real, and thus its kernel $V_{\mathbf{x}}$ is also real.

This implies the Shapiro Conjecture, as spaces $V_{\mathbf{x}}$ for \mathbf{x} a critical point give all spaces of polynomials whose Wronskian has roots s .

Thank you for your attention!

