



Semialgebraic Splines

Approximation Theory 2016 San Antonio
Approximation Theory and Algebraic Geometry
22 May 2016



Frank Sottile

sottile@math.tamu.edu



Work with Michael DiPasquale and Lanyin Sun

Motivating Goals


I: Compute dimensions of spaces of splines on a semi-algebraic cell decomposition, to illustrate some phenomena not observed in traditional splines on simplicial or polyhedral complexes.

II: Learn something about splines and algebraic geometry.

Definition: A (*basic*) *semialgebraic set* is one of the form

$$\{x \in \mathbb{R}^2 \mid h_i(x) \geq 0, \text{ for } i = 1, \dots, m\},$$

where h_1, \dots, h_m are polynomials.



Simplicial Complex

Complex with Semialgebraic Cells

Semialgebraic Splines

A *semialgebraic spline* is a function that is piecewise a polynomial with respect to a complex Δ whose cells are semialgebraic sets.

$C_d^r(\Delta)$: vector space of splines on Δ of degree $\leq d$ and smoothness r .

Consider planar complexes Δ with a single interior vertex, \mathbf{v} , and whose edges are defined by polynomials g_1, \dots, g_N (with $g_i(\mathbf{v}) = 0$), where $\deg(g_i) = n_i$ (In our examples here, $N = 3$ and $n_i = 2$).

Set $S := \mathbb{R}[x, y, z]$ and let $J(\mathbf{v}) := \langle g_i \mid i = 1, \dots, N \rangle$, a homogeneous ideal.

$$\dim C_d^r(\Delta) = \sum_{i=1}^N \binom{d - (r+1)n_i + 2}{2} + \dim(S/J(\mathbf{v}))_d$$

Pencils

Suppose that g_1, \dots, g_N all have degree n and form a pencil (dimension of linear span is 2). Suppose they define s distinct curves. Set

$$t := \min\{s, r + 2\}, \quad a := \lfloor \frac{r+1}{t-1} \rfloor,$$

$$s_1 := (t-1)a + t - r - 2, \quad s_2 := r + 1 - (t-1)a.$$

Theorem. $\dim C_d^r(\Delta)$ equals $\binom{d+2}{2} + (N - t) \binom{d - (r+1)n + 2}{2} + s_1 \binom{d - (r+1+a)n + 2}{2} + s_2 \binom{d - (r+2+a)n + 2}{2}$,

where $\binom{a}{b} = 0$ is zero if $a < b$. For $d > (r+2+a)n + 1$ this is

$$N \binom{d - (r+1)n + 2}{2} + n^2 \left(\binom{a+r+2}{2} - t \binom{a+1}{2} \right).$$


Consequently, the dimension of the spline space does not depend upon any real geometry of the curves underlying the edges. They define n^2 points in the complex projective plane, counted with multiplicity.

Pencils, continued


Theorem. For $d > (r+2+a)n+1$, $\dim C_d^r(\Delta)$ is

$$N \binom{d-(r+1)n+2}{2} + n^2 \left(\binom{a+r+2}{2} - t \binom{a+1}{2} \right).$$


$\dim C_d^r(\Delta)$ is independent of the real geometry of the edge curves.




Four real points



Two real and two complex points



Two double points



A real point and a triple point at infinity

Distinct Tangents

Another extreme is when g_1, \dots, g_N are smooth at v with distinct tangents, given by L_1, \dots, L_n , respectively.

Theorem. For d sufficiently large, $\dim C_d^r(\Delta)$ equals

$$\sum_{i=1}^N \binom{d-(r+1)n_i+2}{2} + \binom{a+r+2}{2} - t \binom{a+2}{2},$$


where $t := \min\{N, r+1\}$ and $a := \lfloor \frac{r+1}{t-1} \rfloor$.

For this, we prove that the ideals $J(v) = \langle g_i^{r+1} \mid i = 1, \dots, N \rangle$ and $\langle L_i^{r+1} \mid i = 1, \dots, N \rangle$ define schemes (supported at v) with the same multiplicity, even though they are not isomorphic when r is large.

\rightsquigarrow Similar to the classical case of edges having distinct slopes.

Possible Extensions

If g_1, \dots, g_N do not form a pencil, but vanish at another point, there are some subtleties.



As $\dim C_d^r(\Delta) = \binom{d-(r+1)^2+2}{2} + \dim(S/J(v))_d$, we consider $\dim(S/J(v))_d$ for d large and $r = 0, 1, 2, 3, 4$.

	0	1	2	3	4
I	3	9	21	36	57
II	3	10	22	38	60
III	3	11	23	40	63